

Model-checking an Epistemic μ -calculus with Synchronous and Perfect Recall Semantics

Rodica Bozianu^{1,3}, Cătălin Dima¹, and Constantin Enea²

¹ LACL, Université Paris Est-Créteil,

61 av. du G-ral de Gaulle, 94010 Créteil, France

² LIAFA, CNRS UMR 7089, Université Paris Diderot - Paris 7,

Case 7014, 75205 Paris Cedex 13, France

³ Faculty of Computer Science, "A.I.Cuza" University
700483 Iași, Romania

Abstract. We show that the model-checking problem is decidable for a fragment of the epistemic μ -calculus with imperfect information and perfect recall. The fragment allows free variables within the scope of epistemic modalities in a restricted form that avoids constructing formulas embodying any form of common knowledge. Our calculus subsumes known decidable fragments of epistemic CTL/LTL , may express winning strategies in two-player games with one player having imperfect information and non-observable objectives, and, with a suitable encoding, decidable instances of the model-checking problem for ATL_{iR} can be encoded as instances of the model-checking problem for the μ -calculus of knowledge.

1 Introduction

The μ -calculus of knowledge is an enrichment of the μ -calculus on trees with individual epistemic modalities K_a (and its dual, denoted P_a). It is designed with the aim that, like the classical modal μ -calculus, it would subsume most combinations of temporal and epistemic logics. The μ -calculus of knowledge is more expressive than linear or branching temporal epistemic logics [12, 20], propositional dynamic epistemic logics [21], or the alternating epistemic μ -calculus [4]. On the other hand, some gaps in its expressive power seem to exist, as witnessed by recent observations in [4] showing that formulas like $\langle\langle a \rangle\rangle p_1 \mathcal{U} p_2$ are not expressible in the fixpoint version of ATL . This expressivity gap can be reproduced in the μ -calculus of knowledge, though the μ -calculus of knowledge is richer than the alternating μ -calculus.

A rather straightforward fragment of the epistemic μ -calculus which has a decidable model-checking problem is the one in which knowledge modalities apply only to closed formulas, that is, formulas in which all second-order variables are bound by some fixpoint operator. The decidability of this fragment follows from recent results on the decidability of the emptiness problem for two player games with one player having incomplete information and with non-observable winning conditions [6].

However more expressive fragments having a decidable model-checking problem seem to exist. For example, winning strategies in two-player games with imperfect information can be encoded as fixpoint formulas in the μ -calculus of knowledge, but not in the above-mentioned restricted fragment. The same holds for some formulas in

ATL with imperfect information and perfect recall (ATL_{iR}) [19, 3]: the ATL formula $\langle\langle a \rangle\rangle \Box p$ can be expressed in a modal μ -calculus of knowledge as $\nu Z. \bigvee_{\alpha \in Act_a} K_a(p \wedge \bigwedge_{\beta \in Act_{Ag \setminus \{a\}}} [\alpha, \beta]Z)$. And there are variants of ATL_{iR} for which the model-checking problem is decidable [7]. Note that a translation of each instance of the model-checking problem for ATL into instances of the model-checking problems for the μ -calculus of knowledge is also possible but requires the modification of the models, as suggested on page 8 below.

Our aim in this paper is to identify such a larger fragment of the epistemic μ -calculus for which model-checking is decidable. The fragment we propose here allows an epistemic modality K_a to be applied to a non-closed μ -calculus formula ϕ , but in such a way that avoids expressing properties that construct any variant of common knowledge for two or more agents. Roughly, the technical restriction is the following: two epistemic operators, referring to the knowledge of two different agents a and b , can be applied to non-closed parts of a formula only if the two agents have *compatible* observations (in the sense that the observability relation of one of the agents is a refinement of the observability relation of the other agent). The variant presented here relies on a *concrete* semantics, in the sense of [9], with the observability relation for each agent a being syntactically identified by a subset Π_a of atomic propositions. We require this in order to syntactically define our fragment of μ -calculus of knowledge with a decidable model-checking problem: the compatibility of two observability relations $\sim_a$ and $\sim_b$ is specified at the syntactic level by imposing that either $\Pi_a \subseteq \Pi_b$ or vice-versa.

The epistemic μ -calculus with perfect recall has a history-based semantics: for each finite transition system T , the formulas of the epistemic μ -calculus must be interpreted over the *tree unfolding* of T . This makes it closer with the tree interpretations of the μ -calculus from [10]. For the classical μ -calculus, there are two ways of proving that the satisfiability and the model-checking problem for the tree interpretation of the logic is decidable: either by providing translations to parity games, or by means of a Finite Model Theorem which ensures that a formula has a tree interpretation iff it has a *state-based* interpretation over a finite transition system (this is known to be equivalent with memoryless determinacy for parity games, see e.g. [5]).

The generalization of the automata approach does not seem to be possible for epistemic μ -calculus, mainly due to the absence of an appropriate generalization of tree automata equivalent with the μ -calculus of knowledge. So we take the approach of providing a generalization of the Finite Model Theorem for our fragment of the epistemic μ -calculus. This result says roughly that the tree interpretation of a formula over the tree unfolding of a given finite transition system T which contains the epistemic operators K_a or P_a is exactly the “tree unfolding” of the finitary interpretation of the formula in a second transition system T' , which is obtained by determinizing the projection of T onto the observations of agent a , a construction that is common for decidable fragments of temporal epistemic logics. Our contribution consists in showing that this construction can be applied for the appropriate fragment of the μ -calculus of knowledge. The proof is given in terms of commutative diagrams between predicate transformers that are the interpretations of non-closed formulas.

The model checking problem for the decidable fragment of the epistemic μ -calculus is non-elementary hard due to the non-elementary hardness of the model-checking problem for the linear temporal logic of knowledge [22]. In the full version of this paper [?], we provide a self-contained proof of this result, by a reduction of the emptiness problem for star-free regular expressions.

The rest of the paper is divided as follows: in the next section we recall the predicate transformer semantics of the μ -calculus and adapt it to our epistemic extension, both for the tree interpretation and the finitary interpretation. We then give our weak variant of the Finite Model Theorem for the classical μ -calculus in the third section. The fourth section serves for introducing our fragment of the epistemic μ -calculus and for proving the decidability of its model-checking problem. We end with a section with conclusions and comments.

2 Preliminaries

We start by fixing a series of notions and notations used in the rest of the paper.

A^* denotes the set of words over A . The length of $\alpha \in A^*$, is denoted $|\alpha|$ and the prefix of α up to position i is denoted $\alpha[1..i]$. Hence, $\alpha[1..0] = \varepsilon$ is the empty word. The (strict) prefix ordering on A^* is denoted $\leq (<)$.

Given a set A and an integer $n \in \mathbb{N}$, an A -tree of *outdegree* $\leq n$ is a partial function $t : [1 \dots n]^* \rightarrow A$ whose support, denoted $\text{supp}(t)$, is a prefix-closed subset of the finite sequences of integers in $[1 \dots n]$. A *node* of t is an element of its support. A *path* in t is a pair (x, ρ) consisting of a node x and the sequence of t -labels of all the nodes which are prefixes of x , $\rho = (t(x[1 \dots i]))_{0 \leq i \leq |x|}$.

A **multi-agent system (MAS, for short)** is a tuple $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi)$ with Ag being the set of agents, Q the set of states, q_0 the initial state of the system, $\delta \subseteq Q \times Q$, Π the set of *atomic propositions*, $\pi : Q \rightarrow 2^\Pi$ and for all $a \in Ag$, $\Pi_a \subseteq \Pi$. A run in the structure M from a state q_0 is an infinite sequence of states $\rho = q_0 q_1 q_2 \dots$ such that $(q_i, q_{i+1}) \in \delta$ for all $i \geq 0$. The set of finite runs in M is denoted $\text{Runs}(M)$. Throughout this paper we consider only finite systems, with $Q = \{1, \dots, n\}$ and $q_0 = 1$, and we assume that Q contains only reachable states.

The 2^Π -tree representing the *unfolding* of a MAS M , denoted t_M , is defined by $\text{supp}(t_M) = \{x \in \mathbb{N}^* \mid 1x \in \text{Runs}(M)\}$ and $t_M(x) = x[|x|]$. For any two positions $x, y \in \text{supp}(t_M)$ with $|x| = |y|$, we denote $x \sim_a y$ if for any $n \leq |x|$ we have that

$$\pi(t(x[1..n])) \cap \Pi_a = \pi(t(y[1..n])) \cap \Pi_a$$

Henceforth, for a word $w \in (2^\Pi)^*$, by $w|_{\Pi_a}$ we denote the sequence defined by $w|_{\Pi_a}[i] = w[i] \cap \Pi_a$ for each $1 \leq i \leq |w|$. Note also that the relation $x \sim_a x'$ is both a relation on the nodes of the tree t_M and on the runs of M .

Predicate transformers: Given a set A , an A -transformer is a mapping $f : (2^A)^n \rightarrow 2^A$.

Following the Knaster-Tarski theorem, any monotone A -transformer $f : 2^A \rightarrow 2^A$ has a unique least and greatest fixpoint, denoted lfp_f , resp. gfp_f .

For an A -transformer $f : (2^A)^n \rightarrow 2^A$, a tuple of sets $B_1, \dots, B_n \subseteq A$ and some $k \leq n$ we denote $f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n) : 2^A \rightarrow 2^A$ the A -transformer with

$$f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n)(B) = f(B_1, \dots, B_{k-1}, B, B_{k+1}, \dots, B_n)$$

Note that when f is monotone, $f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n)$ is monotone too. Hence, both $\text{lfp}_{f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n)}$ and $\text{gfp}_{f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n)}$ exist. These fixpoints can also be seen as the following A -transformers: $\text{lfp}_f^k : (2^A)^n \rightarrow 2^A$ and $\text{gfp}_f^k : (2^A)^n \rightarrow 2^A$, defined respectively as:

$$\begin{aligned} \text{lfp}_f^k(B_1, \dots, B_n) &= \text{lfp}_{f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n)} \\ \text{gfp}_f^k(B_1, \dots, B_n) &= \text{gfp}_{f_k(B_1, \dots, B_{k-1}, \cdot, B_{k+1}, \dots, B_n)} \end{aligned}$$

Note that both these A -transformers are constant in their k -th argument. It is also known that both these A -transformers are monotone if f is monotone.

3 The μ -calculus of Knowledge

Syntax: The syntax of the μ -calculus of knowledge (in positive form) is based on the following sets of symbols: a finite set of *agents* Ag , a family of finite sets of *atomic propositions* $(\Pi_a)_{a \in Ag}$ (no restrictions apply on the pairwise intersections between these sets), with $\Pi = \bigcup_{a \in Ag} \Pi_a$, and a finite set of *second-order variables* $\mathcal{Z} = \{Z_1, \dots, Z_k\}$. The set Π_a represents the set of atoms whose value is *observable* by agent a at each instant (in the sense to be developed further).

The grammar for the formulas of the μ -calculus of knowledge is:

$$\varphi ::= p \mid \neg p \mid Z \mid \varphi \wedge \varphi \mid \varphi \vee \varphi \mid AX\varphi \mid EX\varphi \mid K_a\phi \mid P_a\phi \mid \mu Z.\varphi \mid \nu Z.\varphi$$

where $p \in \Pi$, $a \in Ag$ and $Z \in \mathcal{Z}$. Formulas of the type $K_a\phi$ are read as *agent a knows that ϕ holds*. The dual of K_a , denoted P_a , (and definable as $P_a\phi = \neg K_a\neg\phi$ if negation were allowed), reads as *agent a considers that ϕ is possible*. As usual, for a subset of agents $A \subseteq Ag$ we may denote E_A the “everybody knows” operator, $E_A\phi = \bigwedge_{a \in A} K_a\phi$.

The fragment of the μ -calculus of knowledge which does not involve the knowledge operator K_a (or its dual) is called here the *plain μ -calculus*, or simply the μ -calculus, when there’s no risk of confusion. As usual, we say that a formula ϕ is *closed* if each variable Z in ϕ occurs in the scope of a fixpoint operator for Z .

We will also briefly consider in this paper the *modal μ -calculus of knowledge*, for the sake of comparison with other combinations of temporal and epistemic logics. It has almost the same grammar, but with the nexttime operators EX and AX replaced with modal nexttime operators $\langle \bar{\alpha} \rangle$, resp. $[\bar{\alpha}]$ with $\bar{\alpha}$ representing a tuple of action symbols $\bar{\alpha} = (\alpha_a)_{a \in Ag}$. Note that the modal μ -calculus of knowledge can be translated to the non-modal μ -calculus of knowledge by converting each action name $\alpha \in Act_a$ into an atomic proposition, so the main results of this paper generalize easily to this calculus.

We give two semantics of the μ -calculus of knowledge: a tree semantics and a finitary semantics. The tree semantics is required because we assume that agents have *perfect recall*, and hence they remember all observations made since the system started. The second is necessary for the decision problem. The equivalence between the two semantics on trees generated by MASs, which gives the decidability of the model-checking problem, is a weak form of *memoryless determinacy for tree automata*. We present here both semantics of the μ -calculus of knowledge in a predicate-transformer flavor, more appropriate for stating a number of properties of the logic.

The tree semantics of the μ -calculus of knowledge is given in terms of $2^{I \cup \mathcal{Z}}$ -trees. For a given tree t , each formula ϕ which contains variables $Z_1, \dots, Z_n$ is associated with a $\text{supp}(t)$ -transformer $\|\phi\| : (2^{\text{supp}(t)})^n \rightarrow 2^{\text{supp}(t)}$ by structural induction, as follows:

- The two atoms p and $\neg p$ are interpreted as constant $\text{supp}(t)$ -transformers $\|p\| : (2^{\text{supp}(t)})^n \rightarrow 2^{\text{supp}(t)}$ and $\|\neg p\| : (2^{\text{supp}(t)})^n \rightarrow 2^{\text{supp}(t)}$, defined by the sets $\|p\| = \{x \in \text{supp}(t) \mid p \in \pi(t(x))\}$, resp. $\|\neg p\| = \{x \in \text{supp}(t) \mid p \notin \pi(t(x))\}$.
- Each variable $Z_i \in \mathcal{Z}$ is interpreted as the i -th projection on $(2^{\text{supp}(t)})^n$, that is, as the $\text{supp}(t)$ -transformer $\|Z_i\| : (2^{\text{supp}(t)})^n \rightarrow 2^{\text{supp}(t)}$ with $\|Z_i\|(S_1, \dots, S_n) = S_i, \forall S_1, \dots, S_n \subseteq \text{supp}(t)$.
- $\|\phi_1 \vee \phi_2\| = \|\phi_1\| \cup \|\phi_2\|$ and $\|\phi_1 \wedge \phi_2\| = \|\phi_1\| \cap \|\phi_2\|$.
- Each of the two nexttime operators is mapped to a $\text{supp}(t)$ -transformer, denoted AX , resp. $EX : 2^{\text{supp}(t)} \rightarrow 2^{\text{supp}(t)}$, defined as follows: for each $S \subseteq \text{supp}(t)$,

$$AX(S) = \{x \in \text{supp}(t) \mid \forall i \in \mathbb{N} \text{ if } xi \in \text{supp}(t) \text{ then } xi \in S\}$$

$$EX(S) = \{x \in \text{supp}(t) \mid \exists i \in \mathbb{N} \text{ with } xi \in \text{supp}(t) \text{ and } xi \in S\}$$

Then $\|AX\phi\| = AX \circ \|\phi\|$, and $\|EX\phi\| = EX \circ \|\phi\|$.

- Each pair of epistemic operators is mapped to $\text{supp}(t)$ -transformers K_a , resp. $P_a : 2^{\text{supp}(t)} \rightarrow 2^{\text{supp}(t)}$, defined as follows: for each $S \subseteq \text{supp}(t)$,

$$K_a(S) = \{x \in \text{supp}(t) \mid \forall y \in \text{supp}(t) \text{ with } x \sim_a y \text{ we have } y \in S\}$$

$$P_a(S) = \{x \in \text{supp}(t) \mid \exists y \in \text{supp}(t) \text{ with } x \sim_a y \text{ and } y \in S\}$$

Then $\|K_a\phi\| = K_a \circ \|\phi\|$ and $\|P_a\phi\| = P_a \circ \|\phi\|$.

- For the fixpoint operators we put $\|\mu Z_i.\phi\| = \text{lfp}_{\|\phi\|}^i$ and $\|\nu Z_i.\phi\| = \text{gfp}_{\|\phi\|}^i$.

Note that the two $\text{supp}(t)$ -transformers K_a and P_a are dual and we have that $K_a(S) = P_a(\overline{S})$, with $\overline{\cdot}$ denoting the set complementation. We also denote $t \models \phi$ iff $\varepsilon \in \|\phi\|$.

The following property says that the μ -calculus of knowledge cannot distinguish between isomorphic trees:

Proposition 1. For any two MASs M_1 and M_2 for which there exists some tree isomorphism $\chi : 2^{\text{supp}(t_{M_1})} \rightarrow 2^{\text{supp}(t_{M_2})}$, and for any μ -calculus of knowledge formula ϕ , the following diagram commutes:

$$\begin{array}{ccc} (2^{\text{supp}(t_1)})^n & \xrightarrow{\|\phi\|_{M_1}} & 2^{\text{supp}(t_2)} \\ (\chi)^n \downarrow & & \downarrow \chi \\ (2^{\text{supp}(t_2)})^n & \xrightarrow{\|\phi\|_{M_2}} & 2^{\text{supp}(t_2)} \end{array} \quad (1)$$

Proof. By straightforward structural induction on the formula ϕ .

Let $S_1, \dots, S_n \subseteq 2^{\text{supp}(t_1)}$. We have to prove that $\chi(\|\phi\|_{M_1}(S_1, \dots, S_n)) = \|\phi\|_{M_2}(\chi(S_1), \dots, \chi(S_n))$.

1. For $\phi = p$ we have that

$$\begin{aligned} \chi(\|\phi\|_{M_1}(S_1, \dots, S_n)) &= \chi(\{x \in t_{M_1} \mid \phi \in t_{M_1}(x)\}) \\ &= \{\chi(x) \mid x \in t_{M_1}, \phi \in t_{M_1}(x)\} \\ &= \{y \in t_{M_2} \mid \exists x \in t_{M_1} \text{ with } \phi \in t_{M_1}(x) \text{ s.t. } \chi(x) = y\} \quad \text{since } \chi \text{ is a bijection} \\ &= \{y \in t_{M_2} \mid \phi \in t_{M_2}(y)\} \\ &= \|\phi\|_{M_2}(\chi(S_1), \dots, \chi(S_n)) \end{aligned}$$

The proof is similar for $\phi = \neg p$.

2. For $\phi = Z_i \in \mathcal{Z}$, $\|\phi\|_{M_1}(S_1, \dots, S_n) = S_i$. Then,

$$\chi(\|\phi\|_{M_1}(S_1, \dots, S_n)) = \chi(S_i) = \|\phi\|_{M_2}(\chi(S_1), \dots, \chi(S_n)).$$

3. For $\phi = \phi_1 \vee \phi_2$, we have $\|\phi\| = \|\phi_1\| \cup \|\phi_2\|$. By assuming that the property holds for ϕ_1 and ϕ_2 , we get

$$\begin{aligned} \chi(\|\phi\|_{M_1}(S_1, \dots, S_n)) &= \chi(\|\phi_1\|_{M_1}(S_1, \dots, S_n) \cup \|\phi_2\|_{M_1}(S_1, \dots, S_n)) \\ &= \chi(\|\phi_1\|_{M_1}(S_1, \dots, S_n)) \cup \chi(\|\phi_2\|_{M_1}(S_1, \dots, S_n)) \\ &= \|\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n)) \cup \|\phi_2\|_{M_2}(\chi(S_1), \dots, \chi(S_n)) \\ &= \|\phi_1 \vee \phi_2\|_{M_2}(\chi(S_1), \dots, \chi(S_n)). \end{aligned}$$

We similar proof can be given for $\phi = \phi_1 \wedge \phi_2$.

4. For $\phi = AX\phi_1$, $\|\phi\| = AX \circ \|\phi_1\|$. We have that

$$\begin{aligned} \chi(\|AX\phi_1\|_{M_1}(S_1, \dots, S_n)) &= \chi(AX(\|\phi_1\|_{M_1}(S_1, \dots, S_n))) \\ &= \{ \chi(x) \mid x \in \text{supp}(t_{M_1}) \text{ and } \forall i \in \mathbb{N}, \text{ if } xi \in \text{supp}(t_{M_1}) \\ &\quad \text{then } t_{M_1}(xi) \in \|\phi_1\|_{M_1}(S_1, \dots, S_n) \} \\ &= \{ y \mid \chi^{-1}(y) \in \text{supp}(t_{M_1}) \text{ and } \forall j \in \mathbb{N}, \text{ if } \chi^{-1}(y)\chi^{-1}(j) \in \text{supp}(t_{M_1}), \\ &\quad \text{then } t_{M_1}(\chi^{-1}(y)\chi^{-1}(j)) \in \|\phi_1\|_{M_1}(S_1, \dots, S_n) \} \text{ since } \chi \text{ is bijective} \\ &= \{ y \mid y \in \text{supp}(t_{M_2}) \text{ and } \forall j \in \mathbb{N}, \text{ if } yj \in \text{supp}(t_{M_2}) \\ &\quad \text{then } t_{M_2}(yj) \in \chi(\|\phi_1\|_{M_1}(S_1, \dots, S_n)) \} \\ &= \{ y \in \text{supp}(t_{M_2}) \mid \forall j \in \mathbb{N} \text{ if } yj \in \text{supp}(t_{M_2}) \\ &\quad \text{then } t_{M_2}(yj) \in \|\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n)) \} \\ &= AX(\|\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n))) \\ &= AX \circ \|\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n)) \\ &= \|AX\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n)). \end{aligned}$$

The proof is similar for $\phi = EX\phi_1$.

5. For $\phi = K_a \phi_1$, $\|\phi\| = K_a \circ \|\phi_1\|$. Then,

$$\begin{aligned}
\chi(\|\phi\|_{M_1}(S_1, \dots, S_n)) &= \chi(K_a(\|\phi_1\|_{M_1}(S_1, \dots, S_n))) \\
&= \chi(\{x \in \text{supp}(t_{M_1}) \mid \forall y \in \text{supp}(t_{M_1}) \text{ with } x \sim_a y \\
&\quad \text{we have } y \in \|\phi_1\|_{M_1}(S_1, \dots, S_n)\}) \\
&= \{\chi(x) \in \text{supp}(t_{M_2}) \mid s \in \text{supp}(t_{M_1}) \text{ and } \forall y \in \text{supp}(t_{M_1}) \text{ with } x \sim_a y \\
&\quad \text{we have } y \in \|\phi_1\|_{M_1}(S_1, \dots, S_n)\} \\
&= \{x' \in \text{supp}(t_{M_2}) \mid \forall \chi^{-1}(y') \in \text{supp}(t_{M_1}) \text{ with } \chi^{-1}(x') \sim_a \chi^{-1}(y') \\
&\quad \text{we have } \chi^{-1}(y') \in \|\phi_1\|_{M_1}(S_1, \dots, S_n)\} \\
&= \{x' \in \text{supp}(t_{M_2}) \mid \forall y' \in \text{supp}(t_{M_2}) \text{ with } x' \sim_a y' \\
&\quad \text{we have } y' \in \chi(\|\phi_1\|_{M_1}(S_1, \dots, S_n))\} \\
&= K_a(\chi(\|\phi_1\|_{M_1}(S_1, \dots, S_n))) \\
&= K_a(\|\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n))) \\
&= K_a \circ \|\phi_1\|_{M_2}(\chi(S_1), \dots, \chi(S_n)) \\
&= \|\phi\|_{M_2}(\chi(S_1), \dots, \chi(S_n)).
\end{aligned}$$

A similar proof can be given for $\phi = P_a \phi_1$.

6. For $\phi = \mu Z_i. \phi_1$, $\|\phi\| = \text{lfp}_{\|\phi\|}^i$. Hence,

$$\begin{aligned}
\chi(\|\phi\|_{M_1}(S_1, \dots, S_n)) &= \chi(\text{lfp}_{\|\phi\|_{M_1}}^i(S_1, \dots, S_n)) \\
&= \chi(\text{lfp}_{\|\phi_1\|_{i, M_1}}(S_1, \dots, S_{i-1}, \cdot, S_{i+1}, \dots, S_n)) \\
&= \chi(\min\{S \mid \|\phi_1\|_{i, M_1}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n) = S\}) \\
&= \{\min \chi(S) \mid S \text{ s.t. } \|\phi_1\|_{i, M_1}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n) = S\} \\
&\quad \chi \text{ is monotonous} \\
&= \min\{\chi(S) \mid \chi(\|\phi_1\|_{i, M_1}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n)) = \chi(S)\} \\
&= \text{lfp}_{\chi(\|\phi_1\|_{i, M_1}(S_1, \dots, S_{i-1}, \cdot, S_{i+1}, \dots, S_n))} \quad \text{inductive step} \\
&= \text{lfp}_{\|\phi_1\|_{i, M_2}(\chi(S_1), \dots, \chi(S_{i-1}), \cdot, \chi(S_{i+1}), \dots, \chi(S_n))} \\
&= \text{lfp}_{\|\phi_1\|_{M_2}}^i(\chi(S_1), \dots, \chi(S_n)) \\
&= \|\phi\|_{M_2}(\chi(S_1), \dots, \chi(S_n)).
\end{aligned}$$

The proof is similar for $\phi = \nu Z_i. \phi_1$.

3.1 Comparison with other temporal epistemic frameworks

We discuss the relationship between the μ -calculus of knowledge and other temporal epistemic logics or game models with imperfect information and perfect recall.

As already noted e.g. in [20], the following fixpoint formula defines the common knowledge operator for two agents: $C_{a,b}\phi = \nu Z. (\phi \wedge K_a Z \wedge K_b Z)$.

On the other hand, it's easy to see that the (modal variant of the) μ -calculus of knowledge is more expressive than the alternating epistemic μ -calculus of [4], due to

the possibility to insert knowledge operators “in between” the quantifiers that occur in the semantics of the coalition operators. The relationship with ATL_{iR} is more involved, as we detail in the sequel.

Given a set of agents $A \subseteq Ag$, denote Act_A the cartesian product of the set of action symbols for each agent in A , $Act_A = \times_{a \in A} Act_a$. Then, formulas of the type $\langle\langle A \rangle\rangle \Box p$ can be expressed as the fixpoint formula $\nu Z. \bigvee_{\alpha \in Act_A} K_a(p \wedge \bigwedge_{\beta \in Act_{Ag \setminus \{a\}}} [\alpha, \beta]Z)$.

Formulas containing the until operator cannot be translated into the μ -calculus of knowledge. The reason is similar to the one explained in [4]: in formulas of the type $\langle\langle a \rangle\rangle \Diamond p$ the objective p might not be observable by the agent a , who might only be able to know that, at some given time instance, sometimes in the past, the objective was achieved on all identically observable traces.

Given an ATL_{iR} formula $\phi = \langle\langle a \rangle\rangle p_1 \mathcal{U} p_2$ where p_1 and p_2 are atomic proposition, a MAS M and a finite run ρ in M , the instance of the model-checking problem $M, \rho \models \phi$ can be translated to an instance of the model-checking problem in the modal μ -calculus of knowledge of the following formula:

$$\mu Z. \bigvee_{\alpha \in Act_a} K_a(p_2 \vee past_{p_2} \vee (p_1 \wedge \bigwedge_{\beta \in Act_{Ag \setminus \{a\}}} [\alpha, \beta]Z))$$

and the *modified* system M' , in which are created some copies of the successors of the states s labelled with the atomic proposition p_2 and the corresponding paths. The copies are labelled with the existing atomic propositions in the successor of s to which is added the new atomic proposition $past_{p_2}$. It will label all the states occurring *after* state s carrying a p_2 . This mechanism is similar with the “bookkeeping” employed in the two-player games utilized in [7] for checking whether the same formula ϕ holds at a state of a MAS.

The formalisation of the modification of M is given below: For any multi-agent system $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi, (Act_a)_{a \in Ag})$, we compute the multi-agent system M' such that $M' = (Q', Ag, \delta', q'_0, \Pi, (\Pi_a)_{a \in Ag}, \pi', (Act'_a)_{a \in Ag})$ with $Q' = Q \times \{0, 1\}$, $q'_0 = (q_0, 0)$, $\pi'(q, 0) = \pi(q)$, $\pi'(q, 1) = \pi(q) \cup \{past_{p_2}\}$, $Act'_a = Act_a \times \{0, 1\}$, $\forall a \in Ag$ and the transition relation defined as: For any transition $q \xrightarrow{(\alpha, \beta)} r$ where $\alpha \in Act_a$ and $\beta = \beta_1, \dots, \beta_n$ with $\beta_i \in Act_{Ag \setminus \{a\}}$, in M' we have:

$$\begin{aligned} & - (q, 0) \xrightarrow{((\alpha, 0), \beta)} (r, 0) \\ & - (q, 1) \xrightarrow{((\alpha, x), \beta)} (r, 1), x \in \{0, 1\} \\ & - (q, 0) \xrightarrow{((\alpha, 1), \beta)} (r, 1) \text{ if } p \in \pi(q) \\ & - (q, 0) \xrightarrow{((\alpha, 1), \beta)} (r, 0) \text{ if } p \notin \pi(q) \end{aligned}$$

Given a run $\rho = q_0 \xrightarrow{\alpha_1} q_1 \xrightarrow{\alpha_2} \dots$ we denote q_i by $\rho[i]$, $i = 0, \dots, |\rho|$ and α_{i+1} by $act(\rho, i)$, $i = 0, \dots, |\rho| - 1$. We redefine the tree unfolding for the multi player games as being a partial mapping $t_M = (t_M^{node}, t_M^{edge})$ with $t_M^{node} : \mathbb{N} \rightarrow Q$, $t_M^{node}(x) = \pi(x[[x]])$ and $t_M^{edge} : \mathbb{N} \rightarrow \prod_{a \in Ag} Act_a$, $t_M^{edge}(xi) = \alpha$ if $x \xrightarrow{\alpha} xi$ where $\alpha = (\alpha_1, \dots, \alpha_k) \in \prod_{a \in Ag} Act_a$. In this case, we say that two runs ρ and ρ' are indistinguishable (observationally equivalent) to a coalition A (and note $\rho \sim_A \rho'$) if $|\rho| = |\rho'|$, $act(\rho, i)|_A = act(\rho', i)|_A$ for all $i < |\rho|$, and $\pi_A(\rho[i]) = \pi_A(\rho'[i])$ for all $i \leq |\rho|$.

We define a strategy, as it is defined in [7], σ for a coalition A as any mapping $\sigma : (2^{\Pi_A})^* \rightarrow Act_A$. A strategy σ is compatible with a run $\rho = q_0 \xrightarrow{\alpha_1} q_1 \xrightarrow{\alpha_2} \dots$ if $\sigma(\pi_A(\rho[0]) \dots \pi_A(\rho[i])) = \alpha_{i+1} \Big|_A$ for all $i \leq |\rho|$. If σ is compatible with a run ρ , then it is compatible with any run that is indistinguishable from ρ to A .

We also use $[\alpha]p$ to express the fact that for all the successors xi of x for which $t_M^{edge}(xi) = \alpha$ we have that $p \in \pi(t_M^{node}(xi))$.

In order to prove the equivalence between the two problems, we prove that for any system M and any ATL_{iR} formula $\phi = \langle\langle a \rangle\rangle p_1 \mathcal{U} p_2$, there exists a system M' as defined below and a formula $\phi' = \mu Z. \bigvee_{\alpha \in Act_a} K_a (p_2 \vee past_{p_2} \vee (p_1 \wedge \bigwedge_{\beta \in Act_{Ag \setminus \{a\}}} [\alpha, \beta] Z))$ such that for any run ρ in M and any run $\bar{\rho}$ in M' for which the projection in M is ρ , $M, \rho \models \phi$ if and only if $M', \bar{\rho} \models \phi'$.

This construction can be extended to the whole ATL by structural induction on the formula.

Multi-player games with incomplete information can also be translated into the μ -calculus of knowledge. Recall briefly that a (synchronous) two-player game is a tuple $G = (Q, Ag, (Act_a)_{a \in Ag}, \delta, Q_0, (Obs_a)_{a \in Ag}, (o_a)_{a \in Ag}, par)$ with Q denoting the set of states, $Ag = \{A, B\}$ the set of players, $\delta \subseteq Q \times \prod_{a \in Ag} Act_a \times Q$ denoting the transition relation, $o_a : Q \rightarrow Obs_a$ denoting the observability relation for player a and $par : Q \rightarrow \mathbb{N}$ defining the *parity* of each state.

A player $a \in Ag$ plays by choosing a *feasible strategy*, which is a mapping $\sigma : (Obs_a)^* \rightarrow Act_a$. A strategy for a is *winning* when all the runs that are compatible with that strategy satisfy the property: the maximal parity of a state which occurs infinitely often in the run is even. The winning condition might be non-observable to a , as it might happen that two identically observable states $q_1, q_2 \in Q$ might have different parities.

The set of winning strategies for a player in a multi-player game with imperfect information is then expressible within the μ -calculus of knowledge, similarly to the known encoding of the set of winning strategies in a parity game into the μ -calculus from e.g. [10, 18]. Assuming that the largest parity in Q is even and the atomic proposition p_i holds exactly in all states with parity i , the following μ -calculus of knowledge formula encodes the winning strategies for player a :

$$\nu Z_n \mu Z_{n-1} \dots \mu Z_1. \bigvee_{\alpha \in Act_a} K_a \bigvee_{i \leq n} (p_i \wedge \bigwedge_{\beta \in Act_{Ag \setminus \{a\}}} [\alpha, \beta] Z_i)$$

3.2 The model-checking problem

The model-checking problem for the μ -calculus of knowledge is the problem of deciding, given a MAS M and a closed formula ϕ , whether $t_M \models \phi$.

The undecidability of the model-checking problem for combinations of temporal and epistemic logics based on a synchronous and perfect recall semantics and containing the common knowledge operator [22, 21] implies the following result.

Theorem 1. *The model-checking problem for the μ -calculus of knowledge is undecidable.*

The next two sections are dedicated to finding a fragment of the μ -calculus of knowledge with a decidable model-checking problem.

4 Revisiting the Decidability of the Model-checking Problem for the Tree Semantics of the plain μ -calculus

Given a multi-agent system $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi)$, and an agent $a \in Ag$, we may define the relation $\Gamma_a^M \subseteq Q \times Q$ as follows: $(q, r) \in \Gamma_a^M$ if for any run ρ in M ending in q (i.e. $\rho[\rho] = q$) there exists a run ρ' ending in r with $\rho \sim_a \rho'$. Whenever the MAS M is understood from the context, we use the notation Γ_a instead of Γ_a^M .

We now define a second semantics for the μ -calculus of knowledge, which works on the *set of states* of a MAS M . Each formula ϕ which contains variables $Z_1, \dots, Z_n$ is associated with a Q -transformer $[\phi] : (2^Q)^n \rightarrow 2^Q$, again by structural induction:

- $[p]$ resp. $[\neg p]$ are the constant Q -transformers $[p] = \{q \in Q \mid p \in \pi(q)\}$, resp. $[\neg p] = \{q \in Q \mid p \notin \pi(q)\}$.
- $[Z_i] : (2^Q)^n \rightarrow 2^Q$ is the i -th projection, i.e., given $S_1, \dots, S_n \subseteq Q$, $[Z_i](S_1, \dots, S_n) = S_i$.
- $[\phi_1 \vee \phi_2] = [\phi_1] \cup [\phi_2]$, and $[\phi_1 \wedge \phi_2] = [\phi_1] \cap [\phi_2]$.
- Both nexttime modalities are associated with Q -transformers $AX^f, EX^f : 2^Q \rightarrow 2^Q$ defined as:

$$AX^f(S) = \{q \in Q \mid \forall r \in Q \text{ if } (q, r) \in \delta \text{ then } r \in S\}$$

$$EX^f(S) = \{q \in Q \mid \exists r \in Q \text{ with } (q, r) \in \delta \text{ and } r \in S\}$$

Then $[AX\phi] = AX^f \circ [\phi]$ and, similarly, $[EX\phi] = EX^f \circ [\phi]$,

- Both epistemic operators are associated with Q -transformers $K_a^f, P_a^f : 2^Q \rightarrow 2^Q$ defined as:

$$K_a^f(S) = \overline{\Gamma_a(\overline{S})} = \{q \in Q \mid \forall s \in Q, \text{ if } (s, q) \in \Gamma_a \text{ then } s \in S\}$$

$$P_a^f(S) = \Gamma_a(S) = \{q \in Q \mid \exists s \in S \text{ s.t. } (s, q) \in \Gamma_a\}$$

Then $[P_a\phi] = P_a^f \circ [\phi]$ and $[K_a\phi] = K_a^f \circ [\phi]$.

- $[\mu Z_i.\phi] = \text{lfp}_{[\phi]}^i$ and $[\nu Z_i.\phi] = \text{gfp}_{[\phi]}^i$.

The following result represents a variant of the Finite Model Theorem for μ -calculus and is proved by structural induction on the formula ϕ in [?]:

Theorem 2. *Given a MAS $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi)$ in which $Q = \{1, \dots, n\}$ and $q_0 = 1$, and a (plain) μ -calculus formula ϕ , the following diagram commutes:*

$$\begin{array}{ccc} (2^Q)^n & \xrightarrow{[\phi]} & 2^Q \\ (t_M^{-1})^n \downarrow & & \downarrow t_M^{-1} \\ (2^{\text{supp}(t)})^n & \xrightarrow{\|\phi\|} & 2^{\text{supp}(t)} \end{array} \quad (2)$$

We also say that the diagram 2 holds (or commutes) for the formula ϕ in the system M .

Proof. We proceed by structural induction on the formula ϕ . Note first that the diagram 2 holds for the base cases:

$$\begin{aligned} t_M^{-1}(\llbracket p \rrbracket) &= \llbracket p \rrbracket & t_M^{-1}(\llbracket \neg p \rrbracket) &= \llbracket \neg p \rrbracket \\ t_M^{-1}(\llbracket Z_i \rrbracket(S_1, \dots, S_n)) &= \llbracket Z_i \rrbracket(S_1, \dots, S_n) \text{ for all } S_j \subseteq Q, 1 \leq j \leq n. \end{aligned}$$

The induction step relies on two groups of properties: on one side, the commutativity of t_M^{-1} with finite unions/intersections, and two commutativity diagrams relating t_M^{-1} with the mappings AX/AX^f , resp. EX/EX^f . The second group of properties is represented by two characterizations for the restrictions of $\text{lfp}_{\llbracket \phi \rrbracket}^i$ and $\text{gfp}_{\llbracket \phi \rrbracket}^i$ on t_M -regular sets of nodes of t_M .

The first group of properties is summarized in the following identities:

1. For any two sets $S_1, S_2 \subseteq Q$,

$$\begin{aligned} t_M^{-1}(S_1 \cup S_2) &= t_M^{-1}(S_1) \cup t_M^{-1}(S_2) \\ t_M^{-1}(S_1 \cap S_2) &= t_M^{-1}(S_1) \cap t_M^{-1}(S_2) \end{aligned}$$

2. For any set $S \subseteq Q$,

$$\begin{aligned} t_M^{-1}(AX^f(S)) &= AX(t_M^{-1}(S)) \\ t_M^{-1}(EX^f(S)) &= EX(t_M^{-1}(S)) \end{aligned}$$

The following property is essential for the induction step involving the fixpoint operators:

Claim. Suppose ϕ is a μ -calculus formula for which the commutative diagram 2 holds. Given $\vec{S} \in (2^Q)^n$ with $\vec{S} = (S_1, \dots, S_n)$ and an index $i \leq n$, denote $\hat{\phi}_{\vec{S}}^i$ the function

$$\begin{aligned} \hat{\phi}_{\vec{S}}^i : 2^{\text{supp}(t_M)} &\rightarrow 2^{\text{supp}(t_M)} \\ \hat{\phi}_{\vec{S}}^i(T) &= \llbracket \phi \rrbracket(t_M^{-1}(S_1), \dots, t_M^{-1}(S_{i-1}), T, t_M^{-1}(S_{i+1}), \dots, t_M^{-1}(S_n)) \end{aligned} \quad (3)$$

Also denote $\lceil \hat{\phi}_{\vec{S}}^i \rceil$ the function $\lceil \hat{\phi}_{\vec{S}}^i \rceil : 2^Q \rightarrow 2^Q$ with

$$\lceil \hat{\phi}_{\vec{S}}^i \rceil(R) = \llbracket \phi \rrbracket(S_1, \dots, S_{i-1}, R, S_{i+1}, \dots, S_n)$$

Then $\text{lfp}_{\hat{\phi}_{\vec{S}}^i} = t_M^{-1}(\text{lfp}_{\lceil \hat{\phi}_{\vec{S}}^i \rceil})$ and $\text{gfp}_{\hat{\phi}_{\vec{S}}^i} = t_M^{-1}(\text{gfp}_{\lceil \hat{\phi}_{\vec{S}}^i \rceil})$.

Proof. We may prove by induction on $j \in \mathbb{N}$ that

$$(\hat{\phi}_{\vec{S}}^i)^j(\emptyset) = t_M^{-1}(\lceil \hat{\phi}_{\vec{S}}^i \rceil^j(\emptyset)) \quad (4)$$

where the first empty set is an element of $2^{\text{supp}(t_M)}$ whereas the second is an element of 2^Q .

The base case is straightforward, since, for $j = 0$, Identity 4 reduces to $\emptyset = t_M^{-1}(\emptyset)$.

For the induction step we may use the induction hypothesis about the commutative diagram 2 (applied for producing the third identity below) to conclude that:

$$\begin{aligned}
(\hat{\phi}_{\bar{S}}^i)^{j+1}(\emptyset) &= (\hat{\phi}_{\bar{S}}^i)((\hat{\phi}_{\bar{S}}^i)^j(\emptyset)) \\
&= (\hat{\phi}_{\bar{S}}^i)(t_M^{-1}([\hat{\phi}_{\bar{S}}^i]^j(\emptyset))) \\
&= \|\phi\|(t_M^{-1}(S_1), \dots, t_M^{-1}(S_{i-1}), t_M^{-1}([\hat{\phi}_{\bar{S}}^i]^j(\emptyset)), t_M^{-1}(S_{i+1}), \dots, t_M^{-1}(S_n)) \\
&= t_M^{-1}([\phi](S_1, \dots, S_{i-1}, [\hat{\phi}_{\bar{S}}^i]^j(\emptyset), S_{i+1}, \dots, S_n)) \\
&= t_M^{-1}([\hat{\phi}_{\bar{S}}^i]([\hat{\phi}_{\bar{S}}^i]^j(\emptyset))) \\
&= t_M^{-1}([\hat{\phi}_{\bar{S}}^i]^{j+1}(\emptyset))
\end{aligned}$$

We then need to prove that $([\hat{\phi}_{\bar{S}}^i]^j(\emptyset))_{j \geq 0}$ is an increasing sequence of subsets of Q . To that end, we will prove that $[\hat{\phi}_{\bar{S}}^i](\cdot)$ is monotonously increasing in both arguments. That is, $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'')$ for all $S' \subseteq S'' \in 2^Q$ and for all $\bar{S}_1 = (S_{11}, S_{12}, \dots, S_{1n})$ and $\bar{S}_2 = (S_{21}, S_{22}, \dots, S_{2n})$ with $S_{1k} \subseteq S_{2k}$ for all $1 \leq k \leq n$. This can be proved by induction on the structure of ϕ as follows:

1. For $\phi = p$ or $\phi = \neg p$, the property holds since in this case $[\phi]$ is constant.
2. For $\phi = Z_r$, $[\phi]$ is the r -th projection. If $r = i$, then $[\hat{\phi}_{\bar{S}_1}^i](S') = S' \subseteq S'' = [\hat{\phi}_{\bar{S}_2}^i](S'')$. Otherwise, $[\hat{\phi}_{\bar{S}_1}^i](S') = S_{1r} \subseteq S_{2r} = [\hat{\phi}_{\bar{S}_2}^i](S'')$.
3. For $\phi = \phi_1 \vee \phi_2$, $[\phi] = [\phi_1] \cup [\phi_2]$. Since $[\hat{\phi}_{\bar{S}}^i](S') = [\phi](S_1, \dots, S_{i-1}, S', S_{i+1}, \dots, S_n)$, assuming that the property holds for ϕ_1 and ϕ_2 , $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'')$ and $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'')$. Hence, $[\hat{\phi}_{\bar{S}_1}^i](S') \cup [\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'') \cup [\hat{\phi}_{\bar{S}_2}^i](S'')$ and therefore $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'')$. For $\phi = \phi_1 \wedge \phi_2$ the proof is similar.
4. For $\phi = AX\phi_1$, $[\phi] = AX_f \circ [\phi_1] = \{q \in Q \mid \forall r \in Q, \text{ if } q \rightarrow r \in \delta \text{ then } r \in [\phi_1]\}$. That is, the predecessors of nodes in $[\phi_1]$ that have no successors outside $[\phi_1]$. It is easy to see that AX_f is monotonous using the definition: given $S_1, S_2 \in 2^Q$, $S_1 \subseteq S_2$, we have

$$AX_f(S_1) = \{q \in Q \mid \forall r \in Q, \text{ if } q \rightarrow r \in \delta \text{ then } r \in S_1\}$$

and since $S_1 \subseteq S_2$, if $r \in S_1$, then $r \in S_2$. Hence,

$$AX_f(S_1) \subseteq \{q \in Q \mid \forall r \in Q, \text{ if } q \rightarrow r \in \delta \text{ then } r \in S_2\} = AX_f(S_2).$$

Then, since $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'')$, by applying AX_f which is monotonous, we obtain that $AX_f([\hat{\phi}_{\bar{S}_1}^i](S')) \subseteq AX_f([\hat{\phi}_{\bar{S}_2}^i](S''))$. That is, $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_2}^i](S'')$. A similar proof can be given for $\phi = EX\phi_1$.

5. For $\phi = K_a \phi_1$, $[\phi](S) = K_a^f \circ [\phi_1](S) = \{q \in Q \mid \forall s \in Q \text{ with } (s, q) \in \Gamma_a \text{ then } s \in [\phi_1](S)\}$. We can prove, as we did in the case of AX_f , that K_a^f is monotonous and applying it to $[\hat{\phi}_{1, \bar{S}_1}^i](S') \subseteq [\hat{\phi}_{1, \bar{S}_2}^i](S'')$ from the inductive hypothesis, we obtain that $K_a^f([\hat{\phi}_{1, \bar{S}_1}^i](S')) \subseteq K_a^f([\hat{\phi}_{1, \bar{S}_2}^i](S''))$. That is, $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_1}^i](S')$. For $\phi = P_a \phi_1$ the proof results from the duality of K_a^f and P_a^f , i.e., $P_a^f(S) = \overline{K_a^f(\bar{S})}$. We have that $[\hat{\phi}_{1, \bar{S}_1}^i](S') \subseteq [\hat{\phi}_{1, \bar{S}_2}^i](S'')$ and then $\overline{[\hat{\phi}_{1, \bar{S}_1}^i](S')} \supseteq \overline{[\hat{\phi}_{1, \bar{S}_2}^i](S'')}$.

Applying K_a^f to it and then computing the dual set, we have that $K_a^f(\overline{[\hat{\phi}_{1, \bar{S}_1}^i](S')}) \subseteq K_a^f(\overline{[\hat{\phi}_{1, \bar{S}_2}^i](S'')})$. That is, $P_a^f([\hat{\phi}_{1, \bar{S}_1}^i](S')) \subseteq P_a^f([\hat{\phi}_{1, \bar{S}_2}^i](S''))$.

6. For $\phi = \mu Z_r. \phi_1$, $[\phi] = \text{lfp}_{[\phi_1]}$. We have that

$$\begin{aligned} [\hat{\phi}_{\bar{S}_1}^i](S') &= [\phi](S_{1,1}, \dots, S_{1,i-1}, S', S_{1,i+1}, \dots, S_{1,n}) \\ &= \text{lfp}_{[\phi_1]_r(S_{1,1}, \dots, S_{1,r-1}, \cdot, S_{1,r+1}, \dots, S_{1,i-1}, S', S_{1,i+1}, \dots, S_{1,n})}. \end{aligned}$$

From the inductive hypothesis we have that $[\hat{\phi}_{1, \bar{S}_1}^i](S') \subseteq [\hat{\phi}_{1, \bar{S}_2}^i](S'')$ and then, by applying lfp , we obtain that

$$\begin{aligned} &\text{lfp}_{[\phi_1]_r(S_{1,1}, \dots, S_{1,r-1}, \cdot, S_{1,r+1}, \dots, S_{1,i-1}, S', S_{1,i+1}, \dots, S_{1,n})} \\ &\subseteq \text{lfp}_{[\phi_1]_r(S_{2,1}, \dots, S_{2,r-1}, \cdot, S_{2,r+1}, \dots, S_{2,i-1}, S', S_{2,i+1}, \dots, S_{2,n})} \end{aligned}$$

and then $[\hat{\phi}_{\bar{S}_1}^i](S') \subseteq [\hat{\phi}_{\bar{S}_1}^i](S')$.

We may conclude that $[\hat{\phi}_{\bar{S}}^i]$ is monotonously increasing. Now, we know that $\emptyset \subseteq [\hat{\phi}_{\bar{S}}^i](\emptyset)$.

By induction, we can prove that $[\hat{\phi}_{\bar{S}}^i]^j(\emptyset) \subseteq [\hat{\phi}_{\bar{S}}^i]^{j+1}(\emptyset)$, $\forall j \geq 0$. Hence, $([\hat{\phi}_{\bar{S}}^i]^j(\emptyset))_{j \geq 0}$ is an increasing sequence of subsets.

This sequence stabilizes at a certain integer k , which is the fixpoint of $[\hat{\phi}_{\bar{S}}^i]$:

$$\text{lfp}_{[\hat{\phi}_{\bar{S}}^i]} = [\hat{\phi}_{\bar{S}}^i]^k(\emptyset) = [\hat{\phi}_{\bar{S}}^i]^{k+1}(\emptyset) \quad (5)$$

As a consequence of this and of Identity 4, the fixpoint of $\hat{\phi}_{\bar{S}}^i$ is reached for

$$\text{lfp}_{\hat{\phi}_{\bar{S}}^i} = t_M^{-1}([\hat{\phi}_{\bar{S}}^i]^k(\emptyset))$$

which ends the proof of Claim 4

Returning to the proof of Theorem 2, the induction step concerning the least fixed point follows easily:

$$\begin{aligned} \|\text{lfp}_{\phi}^i\| \circ (t_M^{-1})^n(S_1, \dots, S_n) &= \|\text{lfp}_{\phi}^i\|(t_M^{-1}(S_1), \dots, t_M^{-1}(S_n)) = \text{lfp}_{\|\phi\|(t_M^{-1}(S_1), \dots, t_M^{-1}(S_n))}^i \\ &= \text{lfp}_{\hat{\phi}_{(S_1, \dots, S_n)}^i}^i = t_M^{-1}(\text{lfp}_{[\phi]}^i(S_1, \dots, S_n)) \end{aligned}$$

where in the last step we utilized the claim above. A similar proof gives the commutation property for the greatest fixpoint. $\square$

5 A Fragment of the μ -calculus of Knowledge with a Decidable Model-Checking Problem

In this section, we first introduce some additional notations and notions. Given a MAS M and two agents $a_1, a_2 \in Ag$, we say that the two agents **have compatible observability** if either $\Pi_{a_1} \subseteq \Pi_{a_2}$ or $\Pi_{a_1} \supseteq \Pi_{a_2}$.

Given a formula ϕ , let T_ϕ denote the syntactic tree of ϕ . We also consider that, in T_ϕ , each node labeled with a *variable* also has a *successor*, labeled with $\top$. This convention brings the property that each node in T_ϕ whose formula is a variable has a closed subformula (which is $\top$).

The syntactic tree is constructed by structural induction, with

- $\text{supp}(T_p) = \{\epsilon\}$, $T_p(\epsilon) = p$,
- $\text{supp}(T_{\neg p}) = \{\epsilon\}$, $T_{\neg p}(\epsilon) = \neg p$,
- $\text{supp}(T_Z) = \{\epsilon, 1\}$, $T_Z(\epsilon) = Z$, $T_Z(1) = \top$,
- $\text{supp}(T_{Op\phi_1}) = \{\epsilon\} \cup \{1x \mid x \in \text{supp}(\phi_1)\}$, $T_{Op\phi_1}(\epsilon) = Op$, $T_{Op\phi_1}(1x) = T_{\phi_1}(x)$,
where $Op \in \{AX, EX, K_a, P_a, \mu Z, \nu Z\}$
- $\text{supp}(T_{\phi_1 Op \phi_2}) = \{\epsilon\} \cup \{1x \mid x \in \text{supp}(\phi_1)\} \cup \{2x \mid x \in \text{supp}(\phi_2)\}$, $T_{\phi_1 Op \phi_2}(\epsilon) = Op$, $T_{\phi_1 Op \phi_2}(1x) = T_{\phi_1}(x)$, $T_{\phi_1 Op \phi_2}(2x) = T_{\phi_2}(x)$, $Op \in \{\wedge, \vee\}$

We then denote $\text{form}(x)$ the subformula of ϕ whose syntactic tree is $T_\phi|_x$, i.e. the subtree of T_ϕ rooted at x , and say that x is **closed** if $\text{form}(x)$ is closed.

We then say that an epistemic operator $Op \in \{K_a, P_a \mid a \in Ag\}$ is **non-closed** at a node x in a formula ϕ if $\text{form}(x)$ is not closed, Op labels a node $y \geq x$ and for all the nodes y' lying on the path between x and y we have that $\text{form}(y')$ is not closed.

For each node $x \in \text{supp}(T_\phi)$, we also define $AgNCI_\phi(x)$ as being the set of agents a for which K_a or P_a is non-closed at x . In addition, given two distinct nodes $x_1 < x_2$ with x_2 being closed, we say that x_2 is a *nearest closed successor* of x_1 if no other closed node lies on the path from x_1 to x_2 .

Definition 1. The μ -calculus of non-mixing epistemic fixpoints is the fragment of the μ -calculus of knowledge consisting of formulas ϕ satisfying the following property:

Any two agents a and b for which there exist epistemic operators $Op_a \in \{K_a, P_a\}$, $Op_b \in \{K_b, P_b\}$ such that both Op_a and Op_b are not closed at some node x of T_ϕ must have compatible observability, i.e. $\Pi_a \subseteq \Pi_b$ or $\Pi_b \subseteq \Pi_a$.

All formulas of KB_n [12, 13], that is, CTL with individual knowledge operators, are formulas of the μ -calculus of non-mixing epistemic fixpoints. Other examples of nonmixing formulas are the following (a and b are two agents such that $\Pi_a \subseteq \Pi_b$):

$$\mu Z_1.(p \vee K_a(EX.Z_1) \wedge \nu Z_2.(q \wedge Z_1 \wedge K_a(EX Z_2)))$$

$$\mu Z_1.(p \vee K_a(EX.Z_1) \wedge \nu Z_2.(q \wedge K_b(EX Z_2)))$$

Examples of formulas that are not in the μ -calculus of non-mixing epistemic fixpoints are (a and b are two agents such that $\Pi_a \not\subseteq \Pi_b$ and $\Pi_b \not\subseteq \Pi_a$):

$$C_{a,b}\phi = \nu Z.(\phi \wedge K_a Z \vee K_b Z)$$

$$\mu Z_1.(p \vee K_a(EX.Z_1) \wedge \nu Z_2.(q \wedge Z_1 \wedge K_b(EX Z_2)))$$

Theorem 3. *The model-checking problem for the μ -calculus of non-mixing epistemic fixpoints is decidable.*

The crux of the proof consists of proving a commutativity property relating t_M^{-1} with the operators K_a/K_a^f , resp. P_a/P_a^f , similar with the properties relating t_M^{-1} with AX/AX^f , resp. EX/EX^f . Unfortunately, this commutativity property does not hold for any MAS M , as it is shown by the following example.




Fig. 1. (a) A one-agent system with $\Pi_a = \{p_1\}$, (b) The unfolding of the system in Fig. 1(a)

Example 1. Let M be the one-agent system in Fig. 1. If we put $S = \{1, 3\}$ then using Figure 1 b) and the definitions if K_a/P_a and K_a^f/P_a^f , we have that $K_a^f(\{1, 3\}) = \overline{P_a^f(\{2\})} = \{2, 3\} = \{1\}$. That is,

$$t_M^{-1}(K_a^f(S)) = \{x \in \text{supp}(t) \mid x[x] = 1\}$$

Similarly,

$$\begin{aligned} K_a(t_M^{-1}(\{1, 3\})) &= \overline{P_a(t_M^{-1}(\{2\}))} \\ &= \overline{\{x \in \text{supp}(t) \mid x[x] = 2 \vee (x[x] = 3 \wedge |x| \text{ is even})\}} \\ &= \{x \in \text{supp}(t) \mid x[x] = 1 \vee (x[x] = 3 \wedge |x| \text{ is odd})\}. \end{aligned}$$

We can observe that $t_M^{-1}(K_a^f(S))$ contains only nodes of t_M labeled with state 1, whereas $K_a(t_M^{-1}(S))$ contains more nodes, in particular nodes labeled with 3 occurring on the odd levels of t_M .

Definition 2. Given two MASs $M_i = (Q_i, Ag, \delta_i, q_0^i, \Pi, (\Pi_a)_{a \in Ag}, \pi_i)$ ($i = 1, 2$) over the same set of atomic propositions, we say that M_1 is an **in-splitting** of M_2 if there exists a pair of surjective mappings $\chi = (\chi_{st}, \chi_{tr})$, with $\chi_{st} : Q_1 \rightarrow Q_2$, $\chi_{tr} : \delta_1 \rightarrow \delta_2$ satisfying the following properties:

1. For each $q, r \in Q_1$, $(q, r) \in \delta_1$, $\chi_{tr}((q, r)) = (\chi_{st}(q), \chi_{st}(r)) \in \delta_2$.
2. For each $q \in Q_1$, $\pi_2(\chi_{st}(q)) = \pi_1(q)$.

3. For each $q \in Q_1$, $\text{outdeg}(\chi_{st}(q)) = \text{outdeg}(q)$, where $\text{outdeg}(q)$ is the number of transitions starting in q .
4. $\chi_{st}(q_0^1) = q_0^2$.

The in-splitting is an **isomorphism** whenever χ_{st} and χ_{tr} are bijective.

Further, the pair $\chi = (\chi_{st}, \chi_{tr})$ is called an *in-splitting mapping*. Also, we may write $\chi : M_1 \rightarrow M_2$ to denote the fact that $\chi = (\chi_{st}, \chi_{tr})$ is a witness for M_1 being an in-splitting of M_2 .

Note that an in-splitting mapping (term borrowed from symbolic dynamics [15]) represents a surjective functional bisimulation between two transition systems. The following proposition can be seen as a generalization of this remark (proof given in [?]):

Proposition 2. Consider two MASs $M_i = (Q_i, Ag, \delta_i, q_0^i, \Pi, (\Pi_a)_{a \in Ag}, \pi_i)$ ($i = 1, 2$) over the same set of atomic propositions, connected by an in-splitting mapping $\chi = (\chi_{st}, \chi_{tr}) : M_1 \rightarrow M_2$. Then for any plain μ -calculus formula ϕ the following diagram commutes:

$$\begin{array}{ccc}
 (2^{Q_1})^n & \xrightarrow{[\phi]_{M_1}} & 2^{Q_1} \\
 (\chi_{st}^{-1})^n \uparrow & & \uparrow \chi_{st}^{-1} \\
 (2^{Q_2})^n & \xrightarrow{[\phi]_{M_2}} & 2^{Q_2}
 \end{array} \tag{6}$$

Proof. Let $S_1, \dots, S_n \subseteq 2^{Q_2}$. We prove hence by structural induction on the structure of the formula ϕ that

$$[\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) = \chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n))$$

1. For $\phi = p$ we have

$$\begin{aligned}
 [\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) &= \{q \in Q_1 \mid p \in \pi_1(q)\} \\
 \chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n)) &= \chi_{st}^{-1}(\{q \in Q_2 \mid p \in \pi_2(q)\})
 \end{aligned}$$

Since $\pi_2(\chi_{st}(q)) = \pi_1(q)$ and χ_{st} is surjective, we can conclude that

$$[\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) = \chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n))$$

The proof is similar for $\phi = \neg p$.

2. For $\phi = Z_i$, $[\phi]$ is the i -th projection and hence

$$[\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) = \chi_{st}^{-1}(S_i) = \chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n)).$$

3. For $\phi = \phi_1 \vee \phi_2$, $[\phi] = [\phi_1] \cup [\phi_2]$. Then,

$$\begin{aligned}
 [\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) &= [\phi_1]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) \cup [\phi_2]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) \\
 &= \chi_{st}^{-1}([\phi_1]_{M_2}(S_1, \dots, S_n)) \cup \chi_{st}^{-1}([\phi_2]_{M_2}(S_1, \dots, S_n)) && \text{by induction} \\
 &= \chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n))
 \end{aligned}$$

The proof for $\phi = \phi_1 \wedge \phi_2$ is similar.

4. For $\phi = AX\phi_1$, $[\phi] = AX_f \circ [\phi_1]$.

$$\begin{aligned}
[\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) &= AX_f([\phi_1]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n))) \\
&= \{q \in Q_1 \mid \forall r \in Q_1, \text{ if } q \rightarrow r \in \delta_1 \\
&\quad \text{then } r \in [\phi_1]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n))\} \quad \text{by induction} \\
&= \{q \in Q_1 \mid \forall r \in Q_1, \text{ if } q \rightarrow r \in \delta_1 \text{ then } r \in \chi_{st}^{-1}([\phi_1]_{M_2}(S_1, \dots, S_n))\}
\end{aligned}$$

Further, we want to prove that this set equals to

$$\begin{aligned}
\chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n)) &= \chi_{st}^{-1}(AX_f([\phi_1]_{M_2}(S_1, \dots, S_n))) \\
&= \{\chi_{st}^{-1}(q') \mid q' \in Q_2 \text{ and } \forall r' \in Q_2, \text{ if } q' \rightarrow r' \in \delta_2 \text{ then } r' \in [\phi_1]_{M_2}(S_1, \dots, S_n)\}
\end{aligned}$$

We prove it by double inclusion. Let first take a q in the first set. We have that for all $r \in Q_1$, if $q \rightarrow r \in \delta_1$ then $r \in \chi_{st}^{-1}([\phi_1]_{M_2}(S_1, \dots, S_n))$. From the surjectivity of χ_{st} and χ_{tr} and properties 1) and 3) in Definition 13, there is a $q' \in Q_2$ such that $q \in \chi_{st}^{-1}(q')$ and for all $r' = \chi_{st}(r) \in Q_2$, if $q' \rightarrow r'$ then $r' \in \chi_{st}(\chi_{st}^{-1}([\phi_1]_{M_2}(S_1, \dots, S_n)))$. That is, using again the surjectivity, for all $r' \in Q_2$, if $q' \rightarrow r'$ then $r' \in [\phi_1]_{M_2}(S_1, \dots, S_n)$.

For the inverse inclusion, take $q' \in Q_2$ s.t. for all $r' \in Q_2$, if $q' \rightarrow r' \in \delta_2$ then $r' \in [\phi_1]_{M_2}(S_1, \dots, S_n)$. Again, from the surjectivity of χ_{st} and χ_{tr} and properties 1) and 3) in the above definition we have that there exists $q \in Q_1$ s.t. $\chi_{st}(q) = q'$ and for any transition $q' \rightarrow r' \in \delta_2$ we have transition $q \rightarrow r \in \delta_1$ such that $q' \rightarrow r' = \chi_{st}(q) \rightarrow \chi_{st}(r)$. Since χ_{st} is surjective and property 3) holds, if $q' \in Q_2$, $r' \in Q_2$ with $q' \rightarrow r' \in \delta_2$ implies that $r' \in [\phi_1]_{M_2}(S_1, \dots, S_n)$, then there exists $q \in Q_1$, $q \in \chi_{st}^{-1}(q')$ s.t. for all $r = \chi_{st}^{-1}(r') \in Q_1$, if $q \rightarrow r \in \delta_1$ then $r \in \chi_{st}^{-1}([\phi_1]_{M_2}(S_1, \dots, S_n))$. We can then conclude that $[\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n)) = \chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n))$.

5. For $\phi = \mu Z_r. \phi_1$, $[\phi] = \text{lfp}_{[\phi_1]}^i$.

$$\begin{aligned}
\chi_{st}^{-1}([\phi]_{M_2}(S_1, \dots, S_n)) &= \chi_{st}^{-1}(\text{lfp}_{[\phi_1]_{M_2}}^i(S_1, \dots, S_n)) \\
&= \chi_{st}^{-1}(\text{lfp}_{[\phi]_{i, M_2}}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n)) \\
&= \chi_{st}^{-1}(\min\{S \mid [\phi_1]_{i, M_2}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n) = S\}) \\
&\quad \text{since } \chi_{st}^{-1} \text{ is monotonous} \\
&= \min\{\chi_{st}^{-1}(S) \mid \chi_{st}^{-1}([\phi_1]_{1, M_2}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n)) = \chi_{st}^{-1}(S)\} \\
&= \text{lfp}_{\chi_{st}^{-1}([\phi_1]_{i, M_2}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n))} \quad \text{from inductive hypothesis} \\
&= \text{lfp}_{[\phi_1]_{i, M_1}(\chi_{st}^{-1}(S_1, \dots, S_{i-1}, S, S_{i+1}, \dots, S_n))} \\
&= \text{lfp}_{[\phi_1]_{M_1}}^i(\chi_{st}^{-1}(S_1, \dots, S_n)) \\
&= [\phi]_{M_1}(\chi_{st}^{-1}(S_1, \dots, S_n))
\end{aligned}$$

Remark 4 Proposition 2 does not hold for general μ -calculus of knowledge formulas. To see this, consider the system depicted in Fig. 2 (a), which is an in-splitting of the system from Fig. 1 (a), resulting from splitting state 3 in two states, denoted 3 and 4, (i.e. $\chi(1) = 1, \chi(2) = 2, \chi(3) = \chi(4) = 3$) with transitions $(3, 4) \in \delta$ and $(4, 4) \in \delta$.




Fig. 2. (a) An in-splitting of the system from Fig. 1 ; (b) The unfolding of the system from Fig. 2(a)

Note that $K_a^f(\{1, 4\}) = \overline{P_a^f(\{2, 3\})} = \overline{\{2, 3\}} = \{1, 4\}$ and hence,

$$t_M^{-1}(K_a^f(\{1, 4\})) = \{x \in \text{supp}(t) \mid x[x] = 1 \vee x[x] = 4\}$$

On the other hand,

$$\begin{aligned} K_a(t_M^{-1}(\{1, 4\})) &= \overline{P_a(t_M^{-1}(\{2, 3\}))} \\ &= \overline{\{x \in \text{supp}(t) \mid x[x] = 2 \vee x[x] = 3 \vee (x[x] = 4 \wedge |x| \text{ is even})\}} \\ &= \{x \in \text{supp}(t) \mid x[x] = 1 \vee (x[x] = 4 \wedge |x| \text{ is odd})\}. \end{aligned}$$

That is, $t_M^{-1}(K_a^f(S))$ contains all nodes of t_M labelled with states 1 or 4, whereas $K_a(t_M^{-1}(S))$ contains fewer nodes, in particular nodes labelled with 1 and nodes labelled with 4 occurring on the odd levels of t_M .

The following notion corresponds with the “determinization” used for model-checking LTLK/CTLK [22, 8] or solving 2-player parity games with one player having incomplete information [6]:

Definition 3. Given a MAS $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi)$, we define the multi agent system $\Delta_a^{pre}(M) = (\tilde{Q}^{pre}, Ag, \tilde{\delta}, \tilde{q}_0, \Pi, (\Pi_a)_{a \in Ag}, \tilde{\pi})$ as follows:

- $\tilde{Q}^{pre} = \{(s, S) \mid s \in Q, S \subseteq \{q \in Q \mid \pi_a(q) = \pi_a(s)\}\}$ and $\tilde{q}_0 = (q_0, \{q_0\})$.
- $\tilde{\delta}$ is composed of all tuples of the form $((s, S), (r, R))$ where $(s, r) \in \delta$ and $R = \{r' \in Q \mid \pi_a(r') = \pi_a(r) \text{ and } \exists s' \in S \text{ with } (s', r') \in \delta\}$.
- $\tilde{\pi}(s, S) = \pi(s) = \pi(s)$.

The *a-distinction* of M , denoted $\Delta_a(M)$, is the restriction of $\Delta_a^{pre}(M)$ to reachable states, i.e., $\Delta_a(M) = (\tilde{Q}, Ag, \tilde{\delta} \upharpoonright_{\tilde{Q}}, \tilde{q}_0, \Pi, (\Pi_a)_{a \in Ag}, \tilde{\pi} \upharpoonright_{\tilde{Q}})$ where $\tilde{Q} = \{\tilde{s} \in \tilde{Q}^{pre} \mid \tilde{s} \text{ is reachable from } \tilde{q}_0\}$.

Given a run ρ in $\text{Runs}(\Delta_a(M))$, we denote $\rho|_1$ the projection of ρ onto its first component.

Lemma 1. *Given a MAS $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi)$, the following two properties hold:*

1. *For each run $\bar{\rho}$ in $\Delta_a(M)$ ending in (s, S) ,*

$$S = \{r \in Q \mid \exists \rho' \text{ in } M \text{ that ends in } r \text{ with } \rho' \sim_a \bar{\rho}|_1\}$$

2. *For each two runs $\bar{\rho}, \bar{\rho}'$ in $\Delta_a(M)$ with $\bar{\rho}$ ending in $(s, S) \in \tilde{Q}$, if $\bar{\rho}' \sim_a \bar{\rho}$, then there exists $r \in Q$ s.t. $\bar{\rho}'$ ends in (r, S) .*

Proof. We prove the first property by induction on the length of the path ρ . It easy to see that property holds when $\bar{\rho} = \tilde{q}_0 = (q_0, \{q_0\})$. In this case, $\bar{\rho}'$ can be only $\bar{\rho}$.

Suppose now the property holds for any path $\bar{\rho}$ in $\Delta_a(M)$ with $|\bar{\rho}| = n$. Let $\bar{\rho}'$ in $\Delta_a(M)$ with $|\bar{\rho}'| = n + 1$ that ends in (q, S) . Then exists a path $\bar{\rho}'' = ((q_i, S_i))_{1 \leq i \leq n}$ of length n such that $\bar{\rho}' = \bar{\rho}'' \cdot (q, S)$ with $(q_{n+1}, S_{n+1}) = (q, S)$. From the inductive hypothesis, $S_{n-1} = \{r \in Q \mid \exists \rho \text{ in } M \text{ that ends in } r \text{ with } \rho \sim_a \bar{\rho}''|_1\}$.

Since $(q_{n-1}, S_{n-1}) \rightarrow (q, S) \in \delta$, by definition:

$$S = \{r' \in Q \mid \pi_a(r') = \pi_a(q) \text{ and } \exists s' \in S_{n-1} \text{ with } s' \rightarrow r' \in \delta\}$$

From $s' \in S_{n-1}$ we have that exists ρ in M that ends in s' with $\rho \sim_a \bar{\rho}''|_1$. Because $s' \rightarrow r' \in \delta$, $\pi_a(r') = \pi_a(q)$ and $\bar{\rho}' = \bar{\rho}'' \cdot (q, S)$, we can conclude that there exists ρ' in M , $\rho' = \rho \cdot r'$ that ends in r' with $\rho' \sim_a \bar{\rho}'|_1$. That is,

$$S = \{r' \in Q \mid \exists \rho' \text{ in } M \text{ that ends in } r' \text{ with } \rho' \sim_a \bar{\rho}'|_1\}$$

For the second property we also use the induction on the length of paths ρ and ρ' . The basic case is similar as above, since ρ' can only be ρ .

Suppose the property holds for any ρ and any ρ' of length n with $\rho' \sim_a \rho$ and take $\bar{\rho}$ and $\bar{\rho}' \in \text{Runs}(\Delta_a(M))$ of length $n + 1$, with $\bar{\rho}$ that ends in (s, S) and $\bar{\rho}' \sim_a \bar{\rho}$. Denote $\rho = ((q_i, S_i))_{1 \leq i \leq n}$ and $\rho' = ((q'_i, S'_i))_{1 \leq i \leq n}$. Because $\bar{\rho} \sim_a \bar{\rho}'$, we have that $\rho \sim_a \rho'$ and then $(q'_n, S'_n) = (q_n, S_n)$.

From $(q_n, S_n) \rightarrow (s, S)$ we have that

$$S = \{r'' \in Q \mid \pi_a(r'') = \pi_a(s) \text{ and } \exists s'' \in S' \text{ with } s'' \rightarrow r'' \in \delta\}$$

and from $(q'_n, S_n) \rightarrow (r, R)$,

$$R = \{r'' \in Q \mid \pi_a(r'') = \pi_a(r) \text{ and } \exists s'' \in S' \text{ with } s'' \rightarrow r'' \in \delta\}$$

Since $\bar{\rho} \sim_a \bar{\rho}'$, we have that $\pi_a(s) = \pi_a(r)$ and then $S = R$. That is, $\bar{\rho}'$ ends in (r, S) .

Given a MAS $M = (Q, Ag, \delta, q_0, \Pi, (\Pi_a)_{a \in Ag}, \pi)$, and an agent $a \in Ag$, we say that M is ***a-distinguished*** if Γ_a defined on page 10 is a ***congruence relation***, that is, an equivalence relation with the following property:

for any $q, r \in Q$, if $q\Gamma_a r$, $(q, q') \in \delta$, $(r, r') \in \delta$ and $\pi_a(q') = \pi_a(r')$, then $q'\Gamma_a r'$. (7)

Lemma 2. For a MAS $M = (Q, Ag, \delta, q_0, II, (II_a)_{a \in Ag}, \pi)$ and an agent $a \in Ag$ with Γ_a a congruence relation, we have that $s\Gamma_a r$ if and only if there exists ρ and $\rho' \in \text{Runs}(M)$ s.t. ρ ends in s and ρ' ends in r with $\rho \sim_a \rho'$.

Proof. For the direct implication the proof follows from the definition of Γ_a . The proof in the other direction is made by induction on the length of the path ρ .

We define Γ_a^n with $s\Gamma_a^n r$ if and only if for any run ρ in M ending in s with $|\rho| = n$, there exists a run ρ' ending in r with $\rho \sim_a \rho'$.

We show by induction that Γ_n is a congruence. It is easy to see that for the base case, for any ρ , $|\rho| = 1$, there exists $\rho' = q_0 \sim_a \rho$ since $\rho = q_0$.

Suppose that it holds for n and prove for $n + 1$. Take ρ that ends in s , with $|\rho| = n + 1$ for which there exists ρ' ending in r s.t. $\rho \sim_a \rho'$. That means that there exists $\bar{\rho}$ that ends in $s' = \rho[|\rho| - 1]$, with $|\bar{\rho}| = n$ for which there exists $\bar{\rho}'$ ending in $r' = \rho'[|\rho'| - 1]$ s.t. $\bar{\rho} \sim_a \bar{\rho}'$. From the inductive step, we have that $s'\Gamma_a^n r'$. Since $s' \rightarrow s$, $r' \rightarrow r$, $\pi_a(s) = \pi_a(r)$ and Γ_a is a congruence, we can conclude that $s\Gamma_a^{n+1} r$ and then $s\Gamma_a r$.

Lemma 3. Given a MAS M and an agent $a \in Ag$, for any two reachable states (q, S) and (r, R) in $\Delta_a(M)$, $(q, S)\Gamma_a(r, R)$ if and only if $S = R$.

Proof. We use Lemma 1 for the proof.

In the direct sense, if (q, S) is reachable, we have that $S = \{s \in Q \mid \exists \rho' \text{ in } M, \rho' \sim_a \rho|_1\}$. From $(q, S)\Gamma_a(r, R)$, we have that there exists $\bar{\rho}$ in M ending in (r, R) with $\rho \sim_a \bar{\rho}$ and

$$R = \{s' \in Q \mid \exists \rho'' \text{ in } M, \rho'' \sim_a \bar{\rho}|_1\} = \{s' \in Q \mid \exists \rho'' \text{ in } M, \rho'' \sim_a \rho|_1\} = S$$

In the other direction, let take (q, S) and (r, S) two reachable states in $\Delta_a(M)$ then for all ρ that ends in (q, S) we have that

$$S = \{s \in Q \mid \exists \rho' \text{ in } M, \rho' \sim_a \rho|_1\}$$

Since (r, S) is reachable, $r \in S$. Then there exists ρ' in M , ending in r with $\rho' \sim_a \rho|_1$. Then, by the second point of Lemma 1, there exists $\bar{\rho}'$ in $\Delta_a(M)$ with $\bar{\rho}' \sim_a \rho$ which ends the proof.

Proposition 3. 1. For any MAS M , $\Delta_a(M)$ is an in-splitting of M . We denote this in-splitting as $\Delta_{a,M}^{-1} : \Delta_a(M) \rightarrow M$. Whenever the MAS M is clear from the context, we use the notation Δ_a^{-1} instead of $\Delta_{a,M}^{-1}$.
2. For any agent $a \in Ag$ we have that $\Delta_a(M)$ is a -distinguished.

Proof. For the first property, suppose $\Delta_a(M)$ is an in-splitting of M . We define the following mapping $\chi : \Delta_a(M) \rightarrow M$ for any $q', r' \in \tilde{Q}$, $q' = (q, S_1)$ and $r' = (r, S_2)$ as:

$$\chi_{st}(q') = \chi_{st}(q, S_1) = q$$

$$\chi_{tr}(q' \rightarrow r') = q \rightarrow r$$

These two mappings satisfy the properties from Definition 2 since:

$$\begin{aligned}
\chi_{tr}(q' \rightarrow r') &= q \rightarrow r = \chi_{st}(q') \rightarrow \chi_{st}(r') \\
\pi(\chi_{st}(q')) &= \pi(q) = \tilde{\pi}(q') && \text{by definition 3} \\
outdeg(\chi_{st}(q')) &= outdeg(q) = outdeg(q') && \text{by definition 3} \\
\chi_{st}(q'_0) &= \chi_{st}(q_0, q_0) = q_0
\end{aligned}$$

The surjectivity follows from the definition and the assumption that we work only with MAS in which Q contains only reachable states.

For the second property, we have to prove that Γ_a is a congruence relation over $\Delta_a(M)$. To prove the symmetry, take $(q, S)\Gamma_a(r, R)$ in $\Delta_a(M)$. From Lemma 3, we have that $R = S$.

Let now, take any path $\vec{\rho}'$ in $\Delta_a(M)$ ending in (r, S) . From Lemma 1 we have that

$$S = \{q \in Q \mid \exists \rho \text{ in } M, \text{ ending in } q, \rho \sim_a \vec{\rho}'\}$$

Since (q, S) is reachable, $q \in S$ and then, there exists ρ in M , ending in q such that $\rho \sim_a \vec{\rho}'$. That is, there exists $\bar{\rho}$ ending in (q, S') s.t. $\bar{\rho} \sim_a \vec{\rho}'$. And, using the second property of Lemma 1, we get that $S' = S$, i.e.,

$$\text{there exists } \bar{\rho} \text{ ending in } (q, S) \text{ s.t. } \bar{\rho} \sim_a \vec{\rho}'$$

Hence $(r, S)\Gamma_a(q, S)$ which means that Γ_a is symmetric. Reflexivity and transitivity hold trivially.

For proving that Γ_a is a congruence, note first that, from the definition of $\Delta_a(M)$ if $(q, S) \rightarrow (q', S') \in \tilde{\delta}$, $(r, S) \rightarrow (r', S'') \in \tilde{\delta}$ and $\pi(q') = \pi(r')$, then $S' = S''$.

Suppose now that $(q, S)\Gamma_a(r, S)$, $(q, S) \rightarrow (q', S') \in \tilde{\delta}$, $(r, S) \rightarrow (r', S') \in \tilde{\delta}$ and $\pi(q', S') = \pi(r', S')$. Then, (q', S') and (r', S') are reachable, and using Lemma 3 we may conclude that $(q', S')\Gamma_a(r', S')$ which ends on Γ_a is a congruence relation and $\Delta_a(M)$ is a -distinguished.

Proposition 4. *For any MAS M and two agents $a, b \in Ag$ with $\Pi_a \subseteq \Pi_b$, if M is b -distinguished, then $\Delta_a(M)$ is b -distinguished too.*

Proof. For $\Delta_a(M)$ to be b -distinguished, $\Gamma_b^{\Delta_a(M)}$ has to be a congruence relation. To prove the symmetry of $\Gamma_b^{\Delta_a(M)}$, take $(s, S)\Gamma_b^{\Delta_a(M)}(r, R)$. Since $\Pi_a \subseteq \Pi_b$, we have that $(s, S)\Gamma_a(r, R)$ and then, from Lemma 3, $S = R$.

We then also have that for all $\rho \in \Delta_a(M)$ ending in (s, S) , there exists ρ' ending in (r, S) , $\rho \sim_b \rho'$. That is,

$$\exists \bar{\rho} \text{ in } M \text{ ending in } s \text{ for which } \exists \bar{\rho}' \text{ ending in } r, \bar{\rho} \sim_b \bar{\rho}'$$

Since M is b -distinguished, Γ_b^M must be a congruence relation, and using Lemma 2 we have that $s\Gamma_b^M r$.

Since Γ_b^M is congruence, we have that $r\Gamma_b^M s$. That is,

$$\forall \bar{\rho}' \text{ ending in } r, \exists \bar{\rho} \text{ ending in } s \text{ s.t. } \bar{\rho} \sim_b \bar{\rho}'$$

Let ρ' ending in (r, S) . By lemma 1, we have that $S = \{q \in Q \mid \exists \bar{\rho} \text{ in } M \text{ s.t. } \bar{\rho} \sim_a \rho' \mid_1\}$. Also, (s, S) is reachable, hence, $s \in S$, i.e., there exists $\bar{\rho}$ in M s.t. $\bar{\rho} \sim_b \rho' \mid_1$. Therefore, there must exist $\bar{\bar{\rho}}$ in $\Delta_a(M)$ ending in (s, S') s.t. $\bar{\bar{\rho}} \sim_b \rho'$. Then, from the second property of Lemma 1, we have that $\bar{\bar{\rho}}$ ends in (s, S) , and $(r, S) \Gamma_b(s, S)$. That is, Γ_b is symmetric.

Next, for proving that $\Gamma_b^{\Delta_a(M)}$ is a congruence, suppose that $(r, S) \Gamma_b(s, S)$, $r \rightarrow r' \in \delta$, $s \rightarrow s' \in \delta$ and $\pi_b(s') = \pi_b(r')$. From $(r, S) \Gamma_b(s, S)$ and the fact that Γ_b^M is a congruence relation, applying Lemma 2 we have that $r \Gamma_b^M s$ and then $r' \Gamma_b^M s'$. In a similar way we write the proof for the symmetry, we prove that $(r', S') \Gamma_b(s', S')$, where $S' = \{q' \in Q \mid \pi_a(q') = \pi_a(r') \text{ and } \exists q \in S, q \rightarrow q' \in \delta\}$.

The close relationship between the relation Γ_a and the epistemic operators is resumed by the following proposition:

Proposition 5. *For any MAS M , the following diagram commutes iff M is a -distinguished:*

$$\begin{array}{ccc} 2^Q & \xrightarrow{K_a^f} & 2^Q \\ t_M^{-1} \downarrow & & \downarrow t_M^{-1} \\ 2^{\text{supp}(t_M)} & \xrightarrow{K_a} & 2^{\text{supp}(t_M)} \end{array} \quad (8)$$

The same holds if the pair K_a/K_a^f is replaced with P_a/P_a^f .

Proof. Suppose first that the diagram holds and prove that Γ_a is a congruence.

For reflexivity the proof is straightforward and the transitivity results from the transitivity of $\sim_a$.

To prove the symmetry, take $q \Gamma_a r$. From $q \Gamma_a r$ we have that $t_M^{-1}(\{r\}) \subseteq P_a(t_M^{-1}(\{q\}))$.

By the diagram commutativity for P_a/P_a^f , we have that $P_a(t_M^{-1}(\{q\})) = t_M^{-1}(P_a^f(\{q\}))$ and then $t_M^{-1}(\{r\}) \subseteq t_M^{-1}(P_a^f(\{q\}))$. Applying t_M , we obtain $r \in P_a^f(q)$. That is $r \Gamma_a q$. Now, if the diagram commutes for the pair K_a/K_a^f , recall first that for all $S \subseteq 2^{\text{supp}(t_M)}$, $P_a(S) = \overline{K_a(\overline{S})}$ and $P_a^f(S) = \overline{K_a^f(\overline{S})}$, $\forall S \subseteq 2^Q$.

Thereby we have $P_a(t_M^{-1}(S)) = \overline{K_a(t_M^{-1}(\overline{S}))} = \overline{K_a(t_M^{-1}(\overline{S}))} = \overline{t_M^{-1}(K_a^f(\overline{S}))} = t_M^{-1}(\overline{K_a^f(\overline{S})}) = t_M^{-1}(P_a^f(S))$. That is, the diagram commutes for P_a/P_a^f too and we can proceed as above for proving symmetry.

If now the diagram commutes, for proving that Γ_a is a congruence, take $q \Gamma_a r$ and $q \rightarrow q' \in \delta$ and $r \rightarrow r' \in \delta$ and $\pi_a(q') = \pi_a(r')$.

From $q \Gamma_a r$ we have that $r \in P_a^f(q)$ and then $t_M^{-1}(r) \subseteq t_M^{-1}(P_a^f(q)) = P_a(t_M^{-1}(q))$. We get that there exists $x \in \text{supp}(t_M)$, $x[x] = r$ and $x \in P_a(t_M^{-1}(q))$ and therefore there exists $y \in \text{supp}(t_M)$ with $x \sim_a y$ and $y[y] = q$.

Because $x \sim_a y$, $x[x] = r \rightarrow r'$ and $y[y] = q \rightarrow q'$ with $\pi_a(q') = \pi_a(r')$ we obtain that for $x' = xr'$ and $y' = yq'$ with $x' \sim_a y'$ $x' \in P_a(t_M^{-1}(q')) = t_M^{-1}(P_a^f(q'))$. That is, $(q', r') \in \Gamma_a$.

For the inverse implication, suppose that Γ_a is a congruence.

First, we observe that $t_M^{-1}(P_a^f(S)) \subseteq P_a(t_M^{-1}(S))$ holds from the fact that $t_M^{-1}(\Gamma_a(S)) \subseteq P_a(t_M^{-1}(S))$.

Recall that we defined $qI_a^n r$ as: $(q, r) \in I_a^n$ iff $\forall \rho, |\rho| \leq n$ ending in q , $\exists \rho'$ ending in r such that $\rho \sim_a \rho'$.

We also define $P_a^{f,n}(S) = \{q \in S \mid \exists s \in S, (s, q) \in I_a^n\}$.

We now prove by induction that $P_a(t_M^{-1}(r)^{\leq n}) \subseteq t_M^{-1}(P_a^{f,n}(r))$ for all $n \in \mathbb{N}$ and for all $r \in Q$.

For $n = 0$ the inequality trivially holds. Suppose that for n the equation holds for all $q \in Q$. Let $x \in \text{supp}(t_M)$, $|x| = n + 1$, in $P_a(t_M^{-1}(r)^{\leq n+1})$. We want to prove that it is in $t_M^{-1}(P_a^{f,n}(r))$ too.

Since $x \in P_a(t_M^{-1}(r)^{\leq n+1})$, then there exists $y \in \text{supp}(t_M)$ with $x \sim_a y$ and $y[|y|] = r$, $|y| \leq n + 1$.

Let $x = x'i$. Then $x' \in P_a(t_M^{-1}(y[|y| - 1])^{\leq n})$ (from the inductive hypothesis and the fact that $t_M^{-1}(\{r\}) \subseteq t_M^{-1}(P_a^f(\{q\}))$).

From the inductive hypothesis, we obtain that $x' \in t_M^{-1}(P_a^{f,n}(y[|y| - 1]))$ and then $(y[|y| - 1], x'[|x'|]) \in I_a^n$.

But, because $x \sim_a y$, we have that $x'[|x'|] \rightarrow x[|x|] \in \delta$ and $y[|y| - 1] \rightarrow y[|y|] \in \delta$ and $\pi_a(x[|x|]) = \pi_a(y[|y|])$. Then, $y[|y|]I_a x[|x|]$. This means that $x \in t_M^{-1}(P_a^{f,n+1}(r))$.

It results that, for all $n \in \mathbb{N}$ and for all $r \in Q$, $P_a(t_M^{-1}(r)^{\leq n}) \subseteq t_M^{-1}(P_a^{f,n}(r))$ and then $P_a(t_M^{-1}(S)) \subseteq t_M^{-1}(P_a^f(S))$, for all S which ends the proof of the reverse inclusion.

From that and from the first inclusion, we can say that $P_a(t_M^{-1}(S)) = t_M^{-1}(P_a^f(S))$, $\forall S$.

Definition 4. We say that the pair of epistemic operators K_a/K_a^f , resp. P_a/P_a^f , **commutes for M** if the diagram 8 is commutative for the respective pair.

Proposition 5 gives the first restricted form which may lead to the commutativity of Diagram 2 for formulas of the μ -calculus of knowledge. The second restricted form in which the pair K_a/K_a^f (resp. P_a/P_a^f) commutes with a system is stated as point 2 in the next proposition:

Proposition 6. Consider two MASs $M_i = (Q_i, Ag, \delta_i, q_0^i, \Pi, (\Pi_a)_{a \in Ag}, \pi_i)$ ($i = 1, 2$), with $Q_1 = \{1, \dots, n_1\}$ and $Q_2 = \{1, \dots, n_2\}$, related by an in-splitting $\chi = (\chi_{st}, \chi_{tr}) : M_1 \rightarrow M_2$, and define the tree mapping $\hat{\chi} : \text{supp}(t_{M_1}) \rightarrow \text{supp}(t_{M_2})$, where $\hat{\chi}(\varepsilon) = \varepsilon$ and $\hat{\chi}(xi) = \hat{\chi}(x) \cdot \chi_{st}(i)$, for any $x \in \text{supp}(t_{M_1})$ and $i \in Q_1$. Then the following properties hold:

1. $\hat{\chi}$ is a tree isomorphism between t_{M_1} and t_{M_2} and $t_{M_2} \circ \hat{\chi} = \hat{\chi} \circ t_{M_1}$.
2. For any closed formula ϕ of the μ -calculus of knowledge for which the diagram 2 commutes in the system M_2 , the following property holds:

$$\|\phi\|_{M_1} = t_{M_1}^{-1}(\chi_{st}^{-1}(\lceil \phi \rceil_{M_2}))$$

Proof. The first property is implied by the bijectivity of $\hat{\chi}$ and the fact that χ is an in-splitting and the property 3 of Definition 2.

For the second property, we may easily prove that

$$t_{M_2}(\hat{\chi}(x)) = \chi_{st}(t_{M_1}(x)) \quad (9)$$

Let $x' \in \text{supp}(t_{M_1})$. Then, by definition of $\hat{\chi}$, $x' = xi$ for some $x \in \text{supp}(t_{M_1})$ and $i \in Q_1$.

$$t_{M_2}(\hat{\chi}(x')) = t_{M_2}(\hat{\chi}(xi)) = t_{M_2}(\hat{\chi}(x) \cdot \hat{\chi}_{st}(i)) = \hat{\chi}_{st}(i) \quad (10)$$

On the other hand, we have that

$$\chi_{st}(t_{M_1}(x')) = \chi_{st}(t_{M_1}(xi)) = \chi_{st}(i) \quad (11)$$

From the last two equations, we can see that Identity 9 holds for any $x \in \text{supp}(t_{M_1})$.

We may observe that $\hat{\chi}(t_{M_1}(x)) = \hat{\chi}(x[|x|]) = \hat{\chi}(\epsilon) \cdot \chi_{st}(x[|x|]) = \chi_{st}(x[|x|])$.

And from the Identity 9, we have that $\hat{\chi}(t_{M_1}(x)) = t_{M_2}(\hat{\chi}(x))$, for all $x \in \text{supp}(t_{M_1})$.

As a premise of the third property, we may observe that:

$$t_{M_1}^{-1} \circ \chi_{st}^{-1} = \hat{\chi}^{-1} \circ t_{M_2}^{-1}$$

From this, combined with the hypothesis on the commutativity of diagram 2 for ϕ in M_2 , the isomorphism property for $\hat{\chi}$ in Proposition 1, and Identity 9, we get:

$$\|\phi\|_{M_1} = \hat{\chi}^{-1}(\|\phi\|_{M_2}) = \hat{\chi}^{-1}(t_{M_2}^{-1}(\lceil\phi\rceil_{M_2})) = t_{M_1}^{-1}(\chi_{st}^{-1}(\lceil\phi\rceil_{M_2}))$$

Remark 5 The previous proposition tells us that, for closed formulas of the μ -calculus of knowledge for which Diagram 2 commutes in M_2 , in the eventuality that the system M_2 needs to be replaced with a “larger” system M_1 (for reasons related with the “de-terminization” that ensures the first type of commutativity of K_a/P_a), the validity of ϕ on the tree t_{M_1} can be recovered from the set of states $\chi_{st}^{-1}(\lceil\phi\rceil_{M_2})$, through the inverse tree mapping $t_{M_1}^{-1}$.

We have now the essential ingredients that ensure the decidability of the model-checking problem for the μ -calculus of non-mixing epistemic fixpoints. The algorithm runs as follows: we proceed by constructing the state-transformer interpretations of the subformulas of ϕ on the given system M , in a bottom-up traversal of the syntactic tree T_ϕ . As long as we only treat subformulas not containing any epistemic operator, Theorem 2 ensures that these state transformers are correct finitary abstractions of the tree semantics of our subformulas.

The first time we encounter in T_ϕ an epistemic operator K_a/P_a , say, the subformula in the current node is $K_a\phi'$, we need to replace M with its a -distinction, $\Delta_a(M)$, in order for the appropriate diagram to commute. This replacement is easier when ϕ' is a closed plain μ -calculus formula. By combining Propositions 6 and 5, the tree semantics of the formula $K_a\phi'$ can be computed using the state transformer $K_a^f(\Delta_a^{-1}(\lceil\phi'\rceil_M))$ in $\Delta_a(M)$, where $\Delta_a^{-1}(\lceil\phi'\rceil_M)$ represents the set of states in $\Delta_a(M)$ on which ϕ' holds.

The procedure is different when ϕ' is non-closed. In this situation, we cannot determinize M , as observed in the remark 4. Therefore we need to descend along the syntactic tree to *all* the “nearest” nodes whose formulas are closed, and only there apply the a -distinction construction, thanks to Proposition 6.

Suppose even further that ϕ' itself contains other knowledge operators, and some other knowledge operator K_b is encountered during this descent. The “nonmixing” assumption on our formula implies that this other agent b has compatible observability with our a (K_a and K_b are non-closed at the node associated with K_a). Therefore, the a -distinction of the models applied at lower levels commutes with K_b , fact which is ensured by Proposition 5 when the two agents have compatible observability.

This whole process ends when we arrive in the root of the syntactic tree, with an in-splitting M' of the initial system M and a (constant) state-transformer σ , which gives

the finitary abstraction of the set of nodes of the tree t_M where ϕ holds. The following paragraphs formalize this process.

Proof (Proof of Theorem 3). Given a formula ϕ in the μ -calculus of non-mixing epistemic fixpoints and a MAS M , we associate with each node x of T_ϕ an in-splitting mapping, denoted $T_\phi^{Ins}(x)$, such that the following properties hold:

1. For the root ϵ and any not closed node x in $\text{supp}(T_\phi)$, $T_\phi^{Ins}(\epsilon) = id_M$ and $T_\phi^{Ins}(x) = id_{M'}$, with M and M' appropriate MASs.
2. For any $x, xi \in \text{supp}(T_\phi)$, $i \in \{1, 2\}$, $\text{codom}(T_\phi^{Ins}(x)) = \text{dom}(T_\phi^{Ins}(xi))$,
3. For any nodes $x_1, x_2 \in \text{supp}(T_\phi)$ with $x_1 \leq x_2$, the *in-splitting mapping* between the two nodes is the composition of the mappings from x_1 to x_2 . Formally,

$$T_\phi^{Ins}(x_1 \dots x_2) = \text{not } T_\phi^{Ins}(x_1) \circ \dots \circ T_\phi^{Ins}(x_2)$$

Then, for any x_1, x_2 leaves in T_ϕ , $T_\phi^{Ins}(\epsilon \dots x_1) = T_\phi^{Ins}(\epsilon \dots x_2)$, where ϵ is the root.

4. For any node x_1 which is a nearest closed successor of the root ϵ , if $\text{AgNCL}_\phi(\epsilon) = \{a_1, \dots, a_k\}$ and $\Pi_{a_1} \subseteq \dots \subseteq \Pi_{a_k}$, then $T_\phi^{Ins}(x_1)$ has the form:

$$T_\phi^{Ins}(x_1) = \Delta_{a_1}^{-1} \circ \dots \circ \Delta_{a_k}^{-1} \circ \chi, \text{ for some } \chi,$$

Next, assuming that T_ϕ^{Ins} is constructed with all the properties above, we denote $Ins(T_\phi^{Ins}) = T_\phi^{Ins}(\epsilon \dots x)$ with x any leave in T_ϕ .

The construction of T_ϕ^{Ins} proceeds by structural induction on ϕ . Whenever we want to emphasize a property of the root of the syntactic tree T_ϕ , we denote it ϵ^ϕ .

For the base case we put $T_p^{Ins}(\epsilon) = T_{\neg p}^{Ins}(\epsilon) = id_M$, for any $p \in \Pi$. For $\phi = Z$, $Z \in \mathcal{Z}$, note that, by construction, the root of T_Z has a leaf successor which is the only child node. Then, $T_Z^{Ins}(\epsilon) = T_Z^{Ins}(1) = id_M$.

For the induction case, take a formula $\phi = Op.\phi'$ where $Op \in \{AX, EX, \mu Z, \nu Z\}$, and assume $T_{\phi'}^{Ins}(x)$ is defined. Then we put $T_\phi^{Ins}(1x) = T_{\phi'}^{Ins}(x)$ for any node x of $\text{supp}(T_{\phi'})$, and $T_\phi^{Ins}(\epsilon^\phi) = id_{M'}$, where $M' = \text{dom}(T_{\phi'}^{Ins}(\epsilon^{\phi'}))$.

Suppose $\phi = K_a\phi'$ or $\phi = P_a\phi'$. Note that for each node $1x$ which is not closed in T_ϕ , the node x is not closed in $T_{\phi'}$ either. Then we put $T_\phi^{Ins}(1x) = T_{\phi'}^{Ins}(x) = id_{M'}$, with M' the appropriate MAS. We also put $T_\phi^{Ins}(\epsilon^\phi) = Id_{M_0}$ for the appropriate M_0 . Furthermore, for each closed node $1x_1 \in \text{supp}(T_\phi)$ which *not* a nearest closed successor of ϵ^ϕ , we put $T_\phi^{Ins}(1x_1) = T_{\phi'}^{Ins}(x_1)$.

Take further a node $1x_1$ which is a nearest closed successor of the root ϵ^ϕ and $\text{AgNCL}(\epsilon^\phi) = \{a_1, \dots, a_k\}$. By the above property 4 from the induction hypothesis, the in-splitting mapping in x_1 is of the form $T_{\phi'}^{Ins}(x_1) = \Delta_{a_1}^{-1} \circ \dots \circ \Delta_{a_k}^{-1} \circ \chi$ with $\Pi_{a_1} \subseteq \dots \subseteq \Pi_{a_k}$. On the other hand, by the assumption that ϕ is a nonmixing formula, a must have compatible observability with all the agents $a_1, \dots, a_k$. Therefore, there must exist some $i \leq k$ such that $\Pi_{a_1} \subseteq \dots \subseteq \Pi_{a_i} \subseteq \Pi_a \subseteq \Pi_{a_{i+1}} \subseteq \dots \subseteq \Pi_{a_k}$. We then define

$$T_\phi^{Ins}(1x_1) = \Delta_{a_1}^{-1} \circ \dots \circ \Delta_{a_i}^{-1} \circ \Delta_a^{-1} \circ \Delta_{a_{i+1}}^{-1} \circ \dots \circ \Delta_{a_k}^{-1} \circ \chi$$

Note that the domain and the codomain of each $\Delta_{a_j}^{-1}$, ($j \leq i$) are different in T_ϕ^{Ins} from those in $T_{\phi'}^{Ins}$, due to the insertion of Δ_a^{-1} .

According to the above constructions for $\phi = K_a\phi'$ of $\phi = P_a\phi'$, all the four properties are satisfied by T_ϕ^{Ins} , the fourth one resulting from the construction of the in-splitting mapping for the nearest closed successors of the root.

Finally, take $\phi = \phi_1 Op \phi_2$ ($Op \in \{\wedge, \vee\}$). If $T_{\phi_1}^{Ins} = T_{\phi_2}^{Ins}$, put $T_\phi^{Ins}(1n) = T_{\phi_1}^{Ins}(x)$ for all nodes $x \in \text{supp}(T_{\phi_1})$, $T_\phi^{Ins}(2n) = T_{\phi_2}^{Ins}(x)$ for all $n \in \text{supp}(T_{\phi_2})$ and $T_\phi^{Ins}(\epsilon) = Id_M$.

Suppose now $T_{\phi_1}^{Ins} \neq T_{\phi_2}^{Ins}$. Consider $AgNCl(1) = \{a_1, \dots, a_k\}$ and $AgNCl(2) = \{b_1, \dots, b_l\}$ with $\Pi_{a_1} \subseteq \dots \subseteq \Pi_{a_k}$ and $\Pi_{b_1} \subseteq \dots \subseteq \Pi_{b_l}$. Take then a node x_1 which is a nearest closed successor of the root of T_{ϕ_1} , ϵ^{ϕ_1} , and a node x_2 which is a nearest closed successor of ϵ^{ϕ_2} . By the induction hypothesis we have:

$$\begin{aligned} T_\phi^{Ins}(x_1) &= \Delta_{a_1}^{-1} \circ \dots \circ \Delta_{a_k}^{-1} \circ \chi_1 & InS(T_{\phi_1}^{Ins}) &= T_\phi^{Ins}(x_1) \circ \chi'_1 \\ T_\phi^{Ins}(x_2) &= \Delta_{b_1}^{-1} \circ \dots \circ \Delta_{b_l}^{-1} \circ \chi_2 & InS(T_{\phi_2}^{Ins}) &= T_\phi^{Ins}(x_2) \circ \chi'_2 \end{aligned}$$

with appropriate in-splittings $\chi_1, \chi'_1, \chi_2, \chi'_2$.

On the other hand, by the assumption on ϕ being nonmixing, for any $i \leq k, j \leq l$, the two agents a_i and b_j must have compatible observability. It therefore follows that there exists a reordering of the union $\{a_1, \dots, a_k\} \cup \{b_1, \dots, b_l\}$ as $\{c_1, \dots, c_m\}$ such that $\Pi_{c_i} \subseteq \Pi_{c_{i+1}}$ for all $i \leq m-1$. Denote then:

$$\chi_0 = \Delta_{c_1}^{-1} \circ \dots \circ \Delta_{c_m}^{-1}$$

By Proposition 4, χ_0 is a c -distinction for any $c \in \{a_1, \dots, a_k\} \cup \{b_1, \dots, b_l\}$. Also, by property 2 of the induction hypothesis, χ_0 is independent of the choice of the nodes x_1, x_2 .

The same property from the induction hypothesis also ensures that, for any nearest closed successor $\bar{x}_2$ of ϵ^{ϕ_2} , there exist in-splittings $\bar{\chi}_2^{\phi_2, \bar{x}_2}, \tilde{\chi}_2^{\phi_2, \bar{x}_2}$ such that:

$$T_\phi^{Ins}(\bar{x}_2) = \Delta_{b_1}^{-1} \circ \dots \circ \Delta_{b_l}^{-1} \circ \bar{\chi}_2^{\phi_2, \bar{x}_2} \quad InS(T_{\phi_2}^{Ins}) = T_\phi^{Ins}(\bar{x}_2) \circ \tilde{\chi}_2^{\phi_2, \bar{x}_2} \quad (12)$$

We will then construct $T_\phi^{Ins}(\cdot)$ as follows:

1. For each closed node x which is a leaf in T_{ϕ_1} but not a nearest closed successor of ϵ^{ϕ_1} , we put $T_\phi^{Ins}(1x) = T_{\phi_1}^{Ins}(x) \circ \chi_2 \circ \chi'_2$.
2. For each non-leaf, closed node x in T_{ϕ_1} which is not a nearest closed successor of ϵ^{ϕ_1} we copy $T_\phi^{Ins}(1x) = T_{\phi_1}^{Ins}(x)$.
3. For each nearest closed successor x of ϵ^{ϕ_1} which is not a leaf in T_{ϕ_1} we put $T_\phi^{Ins}(1x) = \chi_0 \circ \chi_1$.
4. For each closed node x which is a leaf in T_{ϕ_1} and a nearest closed successor of ϵ^{ϕ_1} , we put $T_\phi^{Ins}(1x) = \chi_0 \circ \chi_1 \circ \chi'_1 \circ \chi_2 \circ \chi'_2$.
5. For each closed node x which is not a close successor of ϵ^{ϕ_2} we copy $T_\phi^{Ins}(2x) = T_{\phi_2}^{Ins}(x)$.
6. For each closed node x which is a nearest closed successor of ϵ^{ϕ_2} we put $T_\phi^{Ins}(2x) = \chi_0 \circ \chi_1 \circ \chi'_1 \circ \bar{\chi}_2^{\phi_2, x}$, where $\bar{\chi}_2^{\phi_2, x}$ is the in-splitting mapping associated to node x as in Identity 12 above.
7. For the root ϵ and the non-closed nodes x of T_ϕ , $T_\phi^{Ins}(\epsilon) = Id_{M'}$ and $T_\phi^{Ins}(x) = Id_{M''}$, with M' and M'' appropriate MASSs.

It's not difficult to see that the resulting mapping $T_{\phi_2}^{Ins}(\cdot)$ satisfies the five desired properties. More specifically, property 2 amounts to the following identity:

$$InS(T_{\phi}^{Ins}) = \chi_0 \circ \chi_1 \circ \chi'_1 \circ \chi_2 \circ \chi'_2$$

Further, let M_x denote the MAS which is the *domain* of the in-splitting $T_{\phi}^{Ins}(x)$, and denote Q_x its state-space. Also, for convenience, we denote $\overline{M}_x$ the MAS which represents the *codomain* of $T_{\phi}^{Ins}(x)$, and $\overline{Q}_x$ its state-space. Note that when $x, x1 \in \text{supp}(T_{\phi})$, $\overline{M}_x = M_{x1}$, and similarly $\overline{M}_x = M_{x2}$ when $x2 \in \text{supp}(T_{\phi})$.

Once we built the tree T_{ϕ}^{Ins} , we associate with each node x in T_{ϕ} a state-transformer that will give all the information on the satisfiability of $form(x)$ in the given model. Formally, we build a tree T_{ϕ}^{str} whose domain is $\text{supp}(T_{\phi}) \setminus \{x \mid T_{\phi}(x) = \perp\}$ and which, for each node x , represents a state-transformer $T_{\phi}^{str}(x) : (2^{Q_x})^n \rightarrow 2^{Q_x}$. The construction will be achieved such that

$$\|form(x)\| \circ (t_{M_x}^{-1})^n = t_{M_x}^{-1} \circ T_{\phi}^{str}(x) \quad (13)$$

for each node x with $form(x) \neq \perp$.

The construction proceeds bottom-up on $\text{supp}(T_{\phi})$. We actually build *two* trees, T_{ϕ}^{str} and $\overline{T}_{\phi}^{str}$, such that $\overline{T}_{\phi}^{str}(x) : (2^{\overline{Q}_x})^n \rightarrow 2^{\overline{Q}_x}$ and $T_{\phi}^{str}(x) = \overline{T}_{\phi}^{str}(x) \circ \left[(T_{\phi}^{Ins}(x))^{-1} \right]^n$, that is,

$$T_{\phi}^{str}(x)(S_1, \dots, S_n) = \overline{T}_{\phi}^{str}(x)((T_{\phi}^{Ins}(x))^{-1}(S_1, \dots, S_n)) \quad (14)$$

Note that, once we build $\overline{T}_{\phi}^{str}(x)$ for a node x , $T_{\phi}^{str}(x)$ is defined by Identity 14, so we only explain the construction for $\overline{T}_{\phi}^{str}(x)$.

For x leave in T_{ϕ} with $T_{\phi}(x) = p \in \Pi$, we put $\overline{T}_{\phi}^{str}(x) = [p]_M$, the constant state-transformer. Recall that we do not define $T_{\phi}^{str}(x)$ for $\overline{T}_{\phi}(x) = \perp$. For $T_{\phi}(x) = Z_i \in \mathcal{Z}$ we put $\overline{T}_{\phi}^{str}(x)(S_1, \dots, S_n) = S_i$, the i -th projection on $(2^{Q_x})^n$.

For nodes x with $T_{\phi}(x) = Op \in \{AX, EX, K_a, P_a \mid a \in Ag\}$ we put

$$\overline{T}_{\phi}^{str}(x)(S_1, \dots, S_n) = Op(T_{\phi}^{str}(x1)(S_1, \dots, S_n))$$

For $T_{\phi}(x) \in \{\wedge, \vee\}$ we put $\overline{T}_{\phi}^{str}(x)(S_1, \dots, S_n) = (T_{\phi}^{str}(x1)(S_1, \dots, S_n))Op(T_{\phi}^{str}(x2)(S_1, \dots, S_n))$.

For $T_{\phi}(x) = \mu Z_i$ with $1 \leq i \leq n$ we put $\overline{T}_{\phi}^{str}(x) = \text{lfp}_{[T_{\phi}^{str}(x1)]}^i$, and, similarly, for $T_{\phi}(x) = \nu Z_i$ we define $\overline{T}_{\phi}^{str}(x) = \text{gfp}_{[T_{\phi}^{str}(x1)]}^i$.

The validity of the Identity 13 is a corollary of Propositions 5 and 6.

The final step consists of checking whether $q_0^{\varepsilon} \in T_{\phi}^{str}(\varepsilon)$, where q_0^{ε} is the initial state in the MAS M_{ε} associated with the root of T_{ϕ} .

The following result follows from a similar result for LTLK from [22]. A self-contained proof can be found in [?]:

Theorem 6. *The model checking problem for the μ -calculus of non-mixing epistemic fixpoints is hard for non-elementary time.*

6 Conclusions and comments

We have presented a fragment of the μ -calculus of knowledge having a decidable model-checking problem. We argued in the introduction that the decidability result does not seem to be achievable using tree automata or multi-player games. Two-player games with one player having incomplete information and with non-observable winning conditions from [6] do not seem to be appropriate for the whole calculus as they are only equivalent with a restricted type of combinations of knowledge operators and fixpoints, as shown on page 9. We conjecture that the formula $\nu Z(p \vee AX.P_a Z)$ is not equivalent with any (tree automaton presentation of a) two-player game with path winning conditions. Translating this formula to a generalized tree automaton seems to require specifying a winning condition on concatenations of finite paths in the tree with “jumps” between two identically-observable positions in the tree. This conjecture extends the non-expressivity results from [4] relating *ATL* and μ -*ATL*.

The second reason for which the above-mentioned generalization would not work comes from results in [9] showing that the satisfiability problem for CTL or LTL is undecidable with the concrete observability relation presented here. It is then expectable that if a class of generalized tree automata is equivalent with the μ -calculus of non-mixing epistemic fixpoints, then that class would have an undecidable emptiness problem and only its “testing problem” would be decidable. Therefore, the classical determinacy argument for two-player games would not be translatable to such a class of automata.

Acknowledgments

Many thanks to D. Guelev for his careful reading of earlier versions of this paper.

References

1. Th. Ågotnes. Action and knowledge in alternating-time temporal logic. *Synthese*, 149(2):375–407, 2006.
2. A. Arnold and D. Niwiński. *Rudiments of μ -calculus*, volume 146 of *Studies in Logic and the Foundations of Mathematics*. North-Holland, 2001.
3. N. Bulling, J. Dix, and W. Jamroga. Model checking logics of strategic ability: Complexity. In M. Dastani, K. V. Hindriks, and J.-J. C. Meyer, editors, *Specification and Verification of Multi-Agent Systems*, pages 125–160. Springer, 2010.
4. N. Bulling and W. Jamroga. Alternating epistemic mu-calculus. In *Proceedings of IJCAI 2011*, pages 109–114. IJCAI/AAAI, 2011.
5. J. Bradfield and C. Stirling. Modal mu-calculi. In J. van Benthem P. Blackburn and F. Wolter, editors, *The Handbook of Modal Logic*, pages 721–756. Elsevier, 2006.
6. K. Chatterjee and L. Doyen. The complexity of partial-observation parity games. In *Proceedings of LPAR-17*, volume 6397 of *Lecture Notes in Computer Science*, pages 1–14. Springer, 2010.
7. C. Dima, C. Enea, and D. Guelev. Model-checking an alternating-time temporal logic with knowledge, imperfect information, perfect recall and communicating coalitions. *Electronic Proceedings in Theoretical Computer Science*, 25:103–117, 2010.
8. C. Dima. Revisiting satisfiability and model-checking for CTLK with synchrony and perfect recall. In *Proceedings of CLIMA IX*, volume 5405 of *LNAI*, pages 117–131, 2008.

9. C. Dima. Non-axiomatizability for linear temporal logic of knowledge with concrete observability. *Journal of Logic and Computation*, pages 939–958, 2011.
10. E. A. Emerson and C. S. Jutla. Tree automata, mu-calculus and determinacy (extended abstract). In *Proceedings of FOCS'91*, pages 368–377. IEEE Computer Society, 1991.
11. V. Goranko and G. van Drimmelen. Complete axiomatization and decidability of alternating-time temporal logic. *TCS*, 353(1-3):93–117, 2006.
12. J. Halpern and M. Vardi. The complexity of reasoning about knowledge and time: Extended abstract. In *Proceedings of STOC'86*, pages 304–315, 1986.
13. J. Halpern and M. Vardi. The complexity of reasoning about knowledge and time. I. Lower bounds. *Journal of Computer System Sciences*, 38(1):195–237, 1989.
14. M. Kacprzak and W. Penczek. Fully symbolic unbounded model checking for alternating-time temporal logic. *Autonomous Agents and Multi-Agent Systems*, 11(1):69–89, 2005.
15. D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, 1995.
16. A. Lomuscio and Fr. Raimondi. Mcmas: A model checker for multi-agent systems. In *Proceedings of TACAS'2006*, volume 3920 of *LNCS*, pages 450–454, 2006.
17. D. Niwiński and I. Walukiewicz. Games for the mu-calculus. *Theor. Comput. Sci.*, 163(1&2):99–116, 1996.
18. J. Obdržálek. *Algorithmic Analysis of Parity Games*. PhD thesis, University of Edinburgh, 2006.
19. P.-Y. Schobbens. Alternating-time logic with imperfect recall. *Electronic Notes in Theoretical Computer Science*, 85(2):82–93, 2004.
20. N. Shilov and N. Garanina. Model checking knowledge and fixpoints. In *Proceedings of FICS'02*, pages 25–39, 2002.
21. J. van Benthem and E. Pacuit. The tree of knowledge in action: Towards a common perspective. In *Proceedings of AiML'06*, pages 87–106. College Publications, 2006.
22. R. van der Meyden and N. Shilov. Model checking knowledge and time in systems with perfect recall (extended abstract). In *Proceedings of FSTTCS'99*, volume 1738 of *LNCS*, pages 432–445, 1999.